arXiv:0901.2432v2 [math.DG] 1 Aug 2009

The mean curvature flow for equifocal submanifolds

Naoyuki Koike

Abstract

In this paper, we investigate the mean curvature flows having an equifocal submanifold in a symmetric space of compact type and its focal submanifolds as initial data. The investigation is performed by investigating the lifts of the submanifolds and the flows to an (infinite dimensional separable) Hilbert space through a Riemannian submersion of the Hilbert space onto the symmetric space.

1 Introduction

The mean curvature flow of a (Riemannian) submanifold $f_0 : M \hookrightarrow N$ is a map $f : M \times [0,\infty) \to N$ such that, for each $t \in [0,T)$, $f_t : M \to N\,(\underset{\rm def}{\Leftrightarrow} f_t(x) = f(x,t)\ (x \in M))$ is an immersion and $f_*((\frac{\partial}{\partial t})_{(x,t)})$ is the mean curvature vector of $f_t : M \hookrightarrow N$, where T is a positive constant or $T = \infty$ and (t) is the natural coordinate of $[0,T)$. Liu-Terng [LT] investigated the mean curvature flow having isoparametric submanifolds (or their focal submanifolds) in a Euclidean space as initial data and obtained the following facts.

Fact 1([LT]). *Let M be a compact isoparametric submanifold in a Euclidean space and C be the Weyl domain of M at $x_0\,(\in M)$. Then the following statements* (i) *and* (ii) *hold:*

(i) *The mean curvature flow M_t having M as initial data converges to a focal submanifold of M in finite time. If a focal map of M onto F is spherical, then the mean curvature flow M_t has type I singularity, that is, $\lim\limits_{t\to T-0}\max_{v\in S^{\perp}M_t}||A^t_v||^2_\infty(T-t) < \infty$, where A^t_v is the shape operator of M_t for v, $||A^t_v||_\infty$ is the sup norm of A^t_v and $S^{\perp}M_t$ is the unit normal bundle of M_t.*

(ii) *For any focal submanifold F of M, there exists a parallel submanifold M' of M such that the mean curvature flow having M' as initial data converges to F in finite time.*

Fact 2([LT]). *Let M and C be as in Fact 1 and σ be a stratum of dimension greater than zero of ∂C. Then the following statements* (i) *and* (ii) *hold:*

(i) *For any focal submanifold F (of M) through σ, the maen curvature flow F_t having F as initial data converges to a focal submanifold F' (of M) through $\partial\sigma$ in finite time. If the fibration of F onto F' is spherical, then the mean curvature flow F_t has type I singularity.*

(ii) For any focal submanifold F (of M) through $\partial\sigma$, there exists a focal submanifold F' (of M) through σ such that the mean curvature flow F'_t having F' as initial data converges to F in finite time.

As a generalized notion of compact isoparametric hypersurfaces in a sphere and a hyperbolic space, and a compact isoparametric submanifolds in a Euclidean space, Terng-Thorbergsson [TT] defined the notion of an equifocal submanifold in a symmetric space as a compact submanifold M satisfying the following three conditions:

(i) the normal holonomy group of M is trivial,

(ii) M has a flat section, that is, for each $x \in M$, $\Sigma_x := \exp^\perp(T^\perp_x M)$ is totally geodesic and the induced metric on Σ_x is flat, where $T^\perp_x M$ is the normal space of M at x and $\exp^\perp$ is the normal exponential map of M.

(iii) for each parallel normal vector field v of M, the focal radii of M along the normal geodesic γ_{v_x} (with $\gamma'_{v_x}(0) = v_x$) are independent of the choice of $x \in M$, where $\gamma'_{v_x}(0)$ is the velocity vector of γ_{v_x} at 0.

On the other hand, Heintze-Liu-Olmos [HLO] defined the notion of an isoparametric submanifold with flat section in a general Riemannian manifold as a submanifold M satisfying the above condition (i) and the following conditions (ii$'$) and (iii$'$):

(ii$'$) for each $x \in M$, there exists a neighborhood U_x of the zero vector (of $T^\perp_x M$) in $T^\perp_x M$ such that $\Sigma_x := \exp^\perp(U_x)$ is totally geodesic and the induced metric on Σ_x is flat,

(iii$'$) sufficiently close parallel submanifolds of M are CMC with respect to the radial direction.

In the case where the ambient space is a symmetric space G/K of compact type, they showed that the notion of an isoparametric submanifold with flat section coincides with that of an equifocal submanifold. The proof was performed by investigating its lift to $H^0([0,1],\mathfrak{g})$ through a Riemannian submersion $\pi \circ \phi$, where π is the natural projection of G onto G/K and ϕ is the parallel transport map for G (which is a Riemannian submersion of $H^0([0,1],\mathfrak{g})$ onto G ($\mathfrak{g}$:the Lie algebra of G)). Let M be an equifocal submanifold in G/K and v be a parallel normal vector field of M. The end-point map $\eta_v(: M \mapsto G/K)$ for v is defined by $\eta_v(x) = \exp^\perp(v_x)$ $(x \in M)$. Set $M_v := \eta_v(M)$. We call M_v a parallel submanifold of M when $\dim M_v = \dim M$ and a focal submanifold of M when $\dim M_v < \dim M$. The parallel submanifolds of M are equifocal. Let $f : M \times [0,T) \to G/K$ be the mean curvature flow having M as initial data. Then, it is shown that, for each $t \in [0,T)$, $f_t : M \hookrightarrow G/K$ is a parallel submanifold of M and hence it is equifocal (see Lemma 3.1). Fix $x_0 \in M$. Let $\widetilde{C}\,(\subset T^\perp_{x_0}M)$ be the fundamental domain containing the zero vector (of $T^\perp_{x_0}M$) of the Coxeter group (which acts on $T^\perp_{x_0}M$) of M at x_0 and set $C := \exp^\perp(\widetilde{C})$, where we note that $\exp^\perp|_{\widetilde{C}}$ is a diffeomorphism onto C. Without loss of generality, we may assume that G is simply connected. Set $\widetilde{M} := (\pi \circ \phi)^{-1}(M)$, which is an isoparametric submanifold in $H^0([0,1],\mathfrak{g})$. Fix $u_0 \in (\pi \circ \phi)^{-1}(x_0)$. The normal space $T^\perp_{x_0}M$ is identified with the normal space $T^\perp_{u_0}\widetilde{M}$ of $\widetilde{M}$ at u_0 through $(\pi \circ \phi)_{*u_0}$. Each parallel submanifold of M intersects with C at the only point and each focal submanifold of M intersects with ∂C at the only point, where ∂C is the boundary of C. Hence, for the mean curvature flow $f : M \times [0,T) \to G/K$ having M as initial data, each $M_t(:= f_t(M))$ intersects with C at the only point. Denote by $x(t)$ this intersection point and define $u : [0,T) \to \widetilde{C}\,(\subset T^\perp_{x_0}M = T^\perp_{u_0}\widetilde{M})$ by $\exp^\perp(u(t)) = x(t)$ $(t \in [0,T))$. Set $\widetilde{M_t} := (\pi \circ \phi)^{-1}(M_t)$ $(t \in [0,T))$. It is shown that $\widetilde{M_t}$ $(t \in [0,T))$ is the mean curvature flow having $\widetilde{M}$ as initial data because the mean curvature vector of $\widetilde{M_t}$ is the horizontal lift of that of M_t through $\pi \circ \phi$. By investigating $u : [0,T) \to T^\perp_{u_0}\widetilde{M}$, we obtain the following fact corresponding to the second-half part of (i) and (ii) of Fact 1.

Theorem A. *Let M be an equifocal submanifold in a symmetric space G/K of compact type and C be the image of the fundamental domain of the Coxeter group of M at $x_0\,(\in M)$ by the normal exponential map. Then the following statements* (i) *and* (ii) *hold:*

(i) *For any focal submanifold F of M, there exists a parallel submanifold M' of M such that the mean curvature flow having M' as initial data converges to F in finite time.*

(ii) *Assume that M is irreducible and the codimension of M is greater than one. If the mean curvature flow M'_t having a parallel submanifold M' of M as initial data converges to a focal submanifold F of M and the fibration of M' onto F is spherical fibration, then the mean curvature flow M'_t has type I singularity.*

Also, we obtain the following fact corresponding to the second-half part of (i) and (ii) of Fact 2 for the mean curvature flow having a focal submanifold of an equifocal submanifold as initial data.

Theorem B. *Let M and C be as in the statement of Theorem A and σ be a stratum of dimension greater than zero of ∂C (which is a stratified space). Then the following statements* (i) *and* (ii) *hold:*

(i) *For any focal submanifold F of M through $\partial\sigma$, there exists a focal submanifold F' of M through σ such that the mean curvature flow having F' as initial data converges to F in finite time.*

(ii) *Assume that M is irreducible and the codimension of M is greater than one. If the mean curvature flow F'_t having a focal submanifold F' of M through σ as initial data converges to a focal submanifold F of M through $\partial\sigma$ and the fibration of F' onto F is spherical fibration, then the mean curvature flow F'_t has type I singularity.*

For an equifocal submanifold in a symmetric space of compact type, the first-half part of the statement (i) of Facts 1 and 2 do not hold. In fact, there exist Hermann actions admitting a minimal principal orbit (which is equifocal) and a minimal singular orbit (which is a focal submanifold of a principal orbit). So the following question arises naturally.

Question. *Let M and σ be as in Theorem B and F be a focal submanifold of M through σ.*

(i) *Does the mean curvature flow having M as initial data converge to a focal submanifold of M in finite time in the case where M is not minimal ?*

(ii) *Does the mean curvature flow having F as initial data converge to a focal submanifold of F through $\partial\sigma$ in finite time in the case where F is not minimal ?*

According to the homogeneity theorem for an equifocal submanifold by Christ [Ch], all irreducible equifocal submanifolds of codimension greater than one in symmetric spaces of compact type are homogeneous. Hence, according to the result by Heintze-Palais-Terng-Thorbergsson [HPTT], they are principal orbits of hyperpolar actions. Furthermore, according to the classification by Kollross [Kol] of hyperpolar actions on irreducible symmetric spaces of compact type, all hyperpolar actions of cohomogeneity greater than one on the symmetric spaces are Hermann actions. Therefore, all equifocal submanifolds of codimension greater than one in irreducible symmetric spaces of compact type are principal orbits of Hermann actions. In the last section, we describe explicitly the mean curvature flows having orbits of some Hermann actions as initial data.

2 Preliminaries

In this section, we briefly review the quantities associated with an isoparametric submanifold in an (infinite dimensional separable) Hilbert space, which was introduced by Terng [T2]. Let M be an isoparametric submanifold in a Hilbert space V.

2.1. Principal curvatures, curvature normals and curvature distributions Let E_0 and E_i $(i \in I)$ be all the curvature distributions of M, where E_0 is defined by $(E_0)_x = \bigcap\limits_{v \in T_x^\perp M} \operatorname{Ker} A_v \, (x \in M)$. For each $x \in M$, we have $T_xM = \overline{(E_0)_x \oplus \left(\bigoplus\limits_{i \in I} (E_i)_x \right)}$, which is the common eigenspace decomposition of A_v's $(v \in T_x^\perp M)$. Also, let λ_i $(i \in I)$ be the principal curvatures of M, that is, λ_i is the section of the dual bundle $(T^\perp M)^*$ of $T^\perp M$ such that $A_v|_{(E_i)_x} = (\lambda_i)_x(v)\mathrm{id}$ holds for any $x \in M$ and any $v \in T_x^\perp M$, and $\mathbf{n}_i$ be the curvature normal corresponding to λ_i, that is, $\lambda_i(\cdot) = \langle \mathbf{n}_i, \cdot \rangle$.

2.2. The Coxeter group associated with an isoparametric submanifold Denote by l_i^x the affine hyperplane $(\lambda_i)_x^{-1}(1)$ in $T_x^\perp M$. The focal set of M at x is equal to the sum $\bigcup\limits_{i \in I} (x + l_i^x)$ of the affine hyperplanes $x + l_i^x$'s $(i \in I)$ in the affine subspace $x + T_x^\perp M$ of V. Each affine hyperplane l_i^x is called a focal hyperplane of M at x. Let W be the group generated by the reflection R_i^x's $(i \in I)$ with respect to l_i^x. This group is independent of the choice x of M up to group isomorphism. This group is called the Coxeter group associated with M. The fundamental domain of the Coxeter group containing the zero vector of $T_x^\perp M$ is given by $\{v \in T_x^\perp M \mid \lambda_i(v) < 1 \, (i \in I)\}$.

2.3. Principal curvatures of parallel submanifolds Let M_v be the parallel submanifold of M for a (non-focal) parallel normal vector field v, that is $M_v = \eta_v(M)$, where η_v is the end-point map for v. Denote by A^v the shape tensor of M_v. This submanifold M_v also is isoparametric and $A^v_w|_{\eta_{v*}(E_i)_x} = \frac{(\lambda_i)_x(w)}{1-(\lambda_i)_x(v_x)}\mathrm{id}$ $(i \in I)$ for any $w \in T^\perp_{\eta_v(x)} M_v$, that is, $\frac{\lambda_i}{1-\lambda_i(v)}$'s $(i \in I)$ are the principal curvatures of M_v and hence $\frac{\mathbf{n}_i}{1-\lambda_i(v)}$'s $(i \in I)$ are the curvature normals of M_v, where we identify $T^\perp_{\eta_v(x)} M_v$ with $T_x^\perp M$.

2.4. The mean curvature vector of a regularizable submanifold Assume that M is regularizable in sense of [HLO], that is, for each normal vector v of M, the regularizable trace $\operatorname{Tr}_r A_v$ and $\operatorname{Tr} A_v^2$ exist, where $\operatorname{Tr}_r A_v$ is defined by $\operatorname{Tr}_r A_v :=$

$\sum_{i=1}^{\infty}(\mu_i^+ + \mu_i^-)$ as $\mathrm{Spec}\,A_v \setminus \{0\} = \{\mu_1^+, \mu_1^-, \mu_2^+, \mu_2^-, \cdots,\}$ $(\mu_1^- < \mu_2^- < \cdots < 0 < \cdots < \mu_2^+ < \mu_1^+)$, where $\mathrm{Spec}\,A_v$ is the spectrum of A_v. Then the mean curvature vector H of M is defined by $\langle H, v\rangle = \mathrm{Tr}_r\,A_v$ $(\forall\, v \in T^\perp M)$.

Let M be an equifocal submanifold in a symmetric space G/K of compact type and set $\widetilde{M} := (\pi\circ\phi)^{-1}(M)$, where π is the natural projection of G onto G/K and $\phi : H^0([0,1],\mathfrak{g}) \to G$ is the parallel transport map for G.

2.5. The mean curvature vector of the lifted submanifold Denote by $\widetilde{H}$ (resp. H) the mean curvature vector of $\widetilde{M}$ (resp. M). Then $\widetilde{M}$ is a regularizable isoparametric submanifold and $\widetilde{H}$ is equal to the horizontal lift of nH^L of nH $(n := \dim M)$ (see Lemma 5.2 of [HLO]).

3 Proofs of Theorems A and B

In this section, we prove Theorem A. Let M be an equifocal submanifold in a symmetric space G/K of compact type, $\pi : G \to G/K$ be the natural projection and ϕ be the parallel transport map for G. Set $\widetilde{M} := (\pi\circ\phi)^{-1}(M)$. Let $\widetilde{C}(\subset T^\perp_{x_0}M)$ be the fundamental domain of the Coxeter group of M at $x_0(\in M)$ containing the zero vector $\mathbf{0}$ of $T^\perp_{x_0}M$ and set $C := \exp^\perp(\widetilde{C})$, where $\exp^\perp$ is the normal exponential map of M. Fix $u_0 \in (\pi\circ\phi)^{-1}(x_0)$. Under the identification of $T^\perp_{x_0}M$ and $T^\perp_{u_0}\widetilde{M}$, the Coxeter group of M is regarded as that of the isoparametric submanifold $\widetilde{M}$. Denote by H (resp. $\widetilde{H}$) the mean curvature vector of M (resp. $\widetilde{M}$). The mean curvature vector H and $\widetilde{H}$ are a parallel normal vector field of M and $\widetilde{M}$, respectively. Let v be a parallel normal vector field of M and v^L be the horizontal lift of v to $H^0([0,1],\mathfrak{g})$, which is a parallel normal vector field of $\widetilde{M}$. Denote by M_v (resp. $\widetilde{M}_{v^L}$) the parallel submanifold $\eta_v(M)$ (resp. $\eta_{v^L}(\widetilde{M})$) of M (resp. $\widetilde{M}$), where η_v (resp. η_{v^L}) is the end-point map for v (resp. v^L). Then we have $\widetilde{M}_{v^L} = (\pi\circ\phi)^{-1}(M_v)$. Denote by H^v (resp. $\widetilde{H}^{v^L}$) the mean curvature vector of M_v (resp. $\widetilde{M}_{v^L}$). Define a vector field X on $\widetilde{C}$ $(\subset T^\perp_{u_0}\widetilde{M} = T^\perp_{x_0}M)$ by $X_w := (\widetilde{H}^{\widetilde{w}})_{u_0+w}$ $(w \in \widetilde{C})$, where $\widetilde{w}$ is the parallel normal vector field of $\widetilde{M}$ with $\widetilde{w}_{u_0} = w$. Let $\xi : (-S,T) \to \widetilde{C}$ be the maximal integral curve of X with $\xi(0) = \mathbf{0}$. Note that S and T are possible be equal to ∞. Let $\widetilde{\xi(t)}$ be the parallel normal vector field of M with $\widetilde{\xi(t)}_{x_0} = \xi(t)$. Let M_t (resp. $\widetilde{M}_t$) be the mean curvature flow having M (resp. $\widetilde{M}$) as initial data.

Lemma 3.1. *For all $t \in [0,T)$, we have $M_t = M_{\widetilde{\xi(t)}}$ and $\widetilde{M}_t = \widetilde{M}_{\widetilde{\xi(t)}^L}$.*

Proof. Fix $t_0 \in [0,T)$. Define a flow $f : \widetilde{M} \times [0,T) \to H^0([0,1],\mathfrak{g})$ by $f(u,t) := \eta_{\widetilde{\xi(t)}^L}(u)$ $((u,t) \in \widetilde{M} \times [0,T))$, where we note that $f_t(\widetilde{M}) = \widetilde{M}_{\widetilde{\xi(t)}^L}$. For simplicity, denote by $\widetilde{H}^{t_0}$ the mean curvature vector of $\widetilde{M}_{\widetilde{\xi(t_0)}^L}$. It is easy to show that $f_*((\frac{\partial}{\partial t})_{(\cdot,t_0)})$ is a parallel normal vector field of $\widetilde{M}_{\widetilde{\xi(t)}^L}$ and that $f_*((\frac{\partial}{\partial t})_{(u_0,t_0)}) = (\widetilde{H}^{t_0})_{f_{t_0}(u_0)}$. On the other hand, since $\widetilde{M}_{\widetilde{\xi(t_0)}^L}$ is isoparametric, $\widetilde{H}^{t_0}$ is also a parallel normal vector field of $\widetilde{M}_{\widetilde{\xi(t_0)}^L}$. Hence we have $f_*((\frac{\partial}{\partial t})_{(\cdot,t_0)}) = \widetilde{H}^{t_0}$. Therefore, it follows from the arbitrariness of t_0 that f is the mean curvature flow having $\widetilde{M}$ as initial data, that is, $\widetilde{M}_{\widetilde{\xi(t)}^L} = \widetilde{M}_t$ $(t \in [0,T))$ holds. Define a flow $\overline{f} : M \times [0,T) \to G/K$ by $\overline{f}(x,t) := \eta_{\widetilde{\xi(t)}}(x)$ $((x,t) \in M \times [0,T))$, where we note that $\overline{f}_t(M) = M_{\widetilde{\xi(t)}}$ $(\overline{f}_t(\cdot) := \overline{f}(\cdot,t))$. For simplicity, denote by H^t the mean curvature vector of $M_{\widetilde{\xi(t)}}$. Fix $t_0 \in [0,T)$. Since $\widetilde{M}_{\widetilde{\xi(t_0)}^L} = (\pi \circ \phi)^{-1}(M_{\widetilde{\xi(t_0)}})$, we have $(H^{t_0})^L = \widetilde{H}^{t_0}$. On the other hand, we have $\overline{f}_*((\frac{\partial}{\partial t})_{(\cdot,t_0)})^L = f_*((\frac{\partial}{\partial t})_{(\cdot,t_0)})(= \widetilde{H}^{t_0})$. Hence we have $\overline{f}_*((\frac{\partial}{\partial t})_{(\cdot,t_0)}) = H^{t_0}$. Therefore, it follows from the arbitrariness of t_0 that $\overline{f}$ is the mean curvature flow having M as initial data, that is, $M_{\widetilde{\xi(t)}} = M_t$ $(t \in [0,T))$. q.e.d.

Clearly we suffice to show the statement of Theorem A in the case where M is full. Hence, in the sequel, we assume that M is full. Denote by Λ the set of all principal curvatures of $\widetilde{M}$. Set $r := \operatorname{codim} M$. It is shown that the set of all focal hyperplanes of $\widetilde{M}$ is given as the sum of finite pieces of infinite parallel families consisting of hyperplanes in $T^{\perp}_{u_0}\widetilde{M}$ which arrange at equal intervals. Let $\{l^{u_0}_{aj} \mid j \in \mathbb{Z}\}$ $(1 \leq a \leq \bar{r})$ be the finite pieces of infinite parallel families consisting of hyperplanes in $T^{\perp}_{u_0}\widetilde{M}$. Since $l^{u_0}_{aj}$'s $(j \in \mathbb{Z})$ arrange at equal intervals, we can express as

$$\Lambda = \mathop{\cup}_{a=1}^{\bar{r}} \{\frac{\lambda_a}{1 + b_a j} \mid j \in \mathbf{Z}\},$$

where λ_a's and b_a's are parallel sections of $(T^{\perp}\widetilde{M})^*$ and positive constants greater than one, respectively, which are defined by $((\lambda_a)_{u_0})^{-1}(1 + b_a j) = l^{u_0}_{aj}$. For simplicity, we set $\lambda_{aj} := \frac{\lambda_a}{1+b_a j}$. Denote by $\mathbf{n}_{aj}$ and E_{aj} the curvature normal and the curvature distribution corresponding to λ_{aj}, respectively. It is shown that, for each a, $\lambda_{a,2j}$'s $(j \in \mathbb{Z})$ have the same multiplicity and so are also $\lambda_{a,2j+1}$'s $(j \in \mathbb{Z})$. Denote by m^e_a and m^o_a the multiplicities of $\lambda_{a,2j}$ and $\lambda_{a,2j+1}$, respectively. Take a parallel normal vector field v of $\widetilde{M}$ with $v_{u_0} \in \widetilde{C}$. Denote by $\widetilde{A}^v$ (resp. $\widetilde{H}^v$) the shape tensor (resp. the mean curvature vector) of the parallel submanifold $\widetilde{M}_v$. Since

$\widetilde{A}^v_w|_{\eta_{v*}(E_{aj})_u} = \frac{(\lambda_{aj})_u(w)}{1-(\lambda_{aj})_u(v_u)}\,\mathrm{id}$ $(w \in T^\perp_u \widetilde{M})$, we have

$$\begin{aligned}\mathrm{Tr}_r \widetilde{A}^v_w &= \sum_{a=1}^{\bar{r}} \left(\sum_{j\in\mathbf{Z}} \frac{m^e_a \lambda_{a,2j}(w)}{1-\lambda_{a,2j}(v_u)} + \sum_{j\in\mathbf{Z}} \frac{m^o_a \lambda_{a,2j+1}(w)}{1-\lambda_{a,2j+1}(v_u)} \right) \\ &= \sum_{a=1}^{\bar{r}} \Bigg((m^e_a + m^o_a) \cot(\frac{\pi}{b_a}(1-\lambda_a(v_u))) \\ &\qquad + (m^e_a - m^o_a) \mathrm{cosec}(\frac{\pi}{b_a}(1-\lambda_a(v_u))) \Bigg) \frac{\pi}{2b_a} \lambda_a(w),\end{aligned}$$

where we use the relation $\frac{\cos\theta+1}{\sin\theta} = \sum\limits_{j\in\mathbf{Z}} \frac{2}{\theta+2j\pi}$. Therefore we have

$$\begin{aligned}\widetilde{H}^v &= \sum_{a=1}^{\bar{r}} \Bigg((m^e_a + m^o_a) \cot(\frac{\pi}{b_a}(1-\lambda_a(v))) \\ &\qquad + (m^e_a - m^o_a) \mathrm{cosec}(\frac{\pi}{b_a}(1-\lambda_a(v))) \Bigg) \frac{\pi}{2b_a} \mathbf{n}_a,\end{aligned} \tag{3.1}$$

where $\mathbf{n}_a := \mathbf{n}_{a0}$.

Now we prove Theorem A.

Proof of Theorem A. In this proof, we use the above notations. Let X be the above vector field on $\widetilde{C}$. From (3.1), we have

$$\begin{aligned}X_w &= \sum_{a=1}^{\bar{r}} \Bigg((m^e_a + m^o_a) \cot(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w))) \\ &\qquad + (m^e_a - m^o_a) \mathrm{cosec}(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w))) \Bigg) \frac{\pi}{2b_a} (\mathbf{n}_a)_{u_0} \\ &= \sum_{a=1}^{\bar{r}} \Bigg(\frac{\cos(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w)))+1}{\sin(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w)))} \times m^e_a \\ &\qquad + \frac{\cos(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w)))-1}{\sin(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w)))} \times m^o_a \Bigg) \frac{\pi}{2b_a} (\mathbf{n}_a)_{u_0}.\end{aligned} \tag{3.2}$$

Denote by $\bar{r}'$ the cardinal number of the set $\{a \,|\, (\lambda_a)^{-1}_{u_0}(1) \cap \partial\widetilde{C}\ :\ \text{open in } \partial\widetilde{C}\}$. Clearly we have $r+1 \le \bar{r}' \le \bar{r}$. By reordering $\{1, \cdots, \bar{r}\}$, we may assume that this set is equal to $\{1, \cdots, \bar{r}'\}$. Fix $w_0 \in \widetilde{C}$ and $a_0 \in \{1, \cdots, \bar{r}'\}$. For simplicity, set $\widetilde{\sigma}_{a_0} := (\lambda_{a_0})^{-1}_{u_0}(1) \cap \partial\widetilde{C}$. Let w'_0 be the point of $\widetilde{\sigma}_{a_0}$ such that $w_0 - w'_0$ is normal to

$\widetilde{\sigma}_{a_0}$ and set $w_0^{\varepsilon} := \varepsilon w_0 + (1-\varepsilon)w_0'$ for $\varepsilon \in (0,1)$. Then we have $\lim\limits_{\varepsilon\to+0}(\lambda_{a_0})_{u_0}(w_0^{\varepsilon}) = 1$ and $\sup\limits_{0<\varepsilon<1}(\lambda_a)_{u_0}(w_0^{\varepsilon}) < 1$ for each $a \in \{1,\cdots,\bar{r}\}\setminus\{a_0\}$. Hence we have

$$\lim_{\varepsilon\to+0}\frac{\cos(\frac{\pi}{b_{a_0}}(1-(\lambda_{a_0})_{u_0}(w_0^{\varepsilon})))+1}{\sin(\frac{\pi}{b_{a_0}}(1-(\lambda_{a_0})_{u_0}(w_0^{\varepsilon})))} = \infty, \quad \lim_{\varepsilon\to+0}\frac{\cos(\frac{\pi}{b_{a_0}}(1-(\lambda_{a_0})_{u_0}(w_0^{\varepsilon})))-1}{\sin(\frac{\pi}{b_{a_0}}(1-(\lambda_{a_0})_{u_0}(w_0^{\varepsilon})))} = 0$$

and

$$\sup_{0<\varepsilon<1}\frac{\cos(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w_0^{\varepsilon})))\pm 1}{\sin(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(w_0^{\varepsilon})))} < \infty.$$

Therefore, we have $\lim\limits_{\varepsilon\to+0}\frac{X_{w_0^{\varepsilon}}}{||X_{w_0^{\varepsilon}}||} = \frac{(\mathbf{n}_{a_0})_{u_0}}{||(\mathbf{n}_{a_0})_{u_0}||}$ and $\lim\limits_{\varepsilon\to+0}||X_{w_0^{\varepsilon}}|| = \infty$. This implies that X is as in Fig. 1 on a sufficiently small collar neighborhood of $\widetilde{\sigma}_{a_0}$. Therefore, the flow of X is as in Fig. 2 on a sufficiently small collar neighborhood of $\partial\widetilde{C}$. Take an arbitrary focal submanifold F of M and set $\widetilde{F} := (\pi\circ\phi)^{-1}(F)$, which is a focal



Fig. 1.



Fig. 2.

submanifold of $\widetilde{M}$ and passes through $\partial\widetilde{C}$. From the above fact for X, we see that there exists a parallel submanifold $\widetilde{M'}$ of $\widetilde{M}$ such that the mean curvature flow $\widetilde{M'_t}$ having $\widetilde{M'}$ as initial data converges to $\widetilde{F}$ in finite time. Then $M' := (\pi\circ\phi)(\widetilde{M'})$ is a parallel submanifold of M. According to Lemma 3.1, $(\pi\circ\phi)(\widetilde{M'_t})$ is the mean curvature flow having M' as initial data and it converges to F in finite time. Thus the statement (i) is shown. Next we shall show the statement (ii). Assume that M is irreducible and the codimension of M is greater than one. Let M' and F be as in the statement (ii). Assume that the mean curvature flow having M' as initial data converges to F in finite time T. Set $\widetilde{M'} := (\pi\circ\phi)^{-1}(M')$ and $\widetilde{F} := (\pi\circ\phi)^{-1}(F)$. Since the fibration of M onto F is a spherical fibration, $\widetilde{F}$ passes through a highest dimensional stratum $\widetilde{\sigma}$ of $\partial\widetilde{C}$. Let a_0 be the element of $\{1,\cdots,\bar{r}'\}$ with $\widetilde{\sigma}\subset(\lambda_{a_0})_{u_0}^{-1}(1)$. Set $\widetilde{M'_t} := (\pi\circ\phi)^{-1}(M'_t)$ $(t\in[0,T))$, which is the mean curvature flow having $\widetilde{M'}$ as initial data. Denote by A^t (resp. $\widetilde{A}^t$) the shape tensor of M'_t (resp. $\widetilde{M'_t}$). Let $\widetilde{M'}\cap(u_0+\widetilde{C})=\{u_1\}$, $\xi:[0,T)\to\widetilde{C}$ be the integral curve of X with $\xi(0)=u_1-u_0$ and $\widetilde{\xi(t)}$ be the parallel normal vector field of M' with $\widetilde{\xi(t)}_{x_0}=\xi(t)$. Then, since $\widetilde{M'_t}$ is the parallel submanifold of $\widetilde{M'}$ for $\widetilde{\xi(t)}^L$, we have

$$\mathrm{Spec}\,\widetilde{A}^t_v\setminus\{0\}=\{\frac{(\lambda_{aj})_{u_0}(v)}{1-(\lambda_{aj})_{u_0}(\xi(t))}\,|\,a=1,\cdots,\bar{r}',\ j\in\mathbb{Z}\}$$

for each $v\in T^{\perp}_{u_0+\xi(t)}\widetilde{M'_t}=T^{\perp}_{u_0}\widetilde{M}$. Since $\lim\limits_{t\to T-0}\xi(t)\in(\lambda_{a_0})_{u_0}^{-1}(1)$, we have $\lim\limits_{t\to T-0}(\lambda_{a_0})_{u_0}(\xi(t))=1$ and $\lim\limits_{t\to T-0}(\lambda_a)_{u_0}(\xi(t))<1$ $(a\neq a_0)$. Hence we have

$$\begin{aligned}&\lim_{t\to T-0}||\widetilde{A}^t_v||^2_\infty(T-t)\\ =&\lim_{t\to T-0}\frac{(\lambda_{a_0})_{u_0}(v)^2}{(1-(\lambda_{a_0})_{u_0}(\xi(t)))^2}(T-t)\\ =&\frac{1}{2}(\lambda_{a_0})_{u_0}(v)^2\lim_{t\to T-0}\frac{1}{(1-(\lambda_{a_0})_{u_0}(\xi(t)))(\lambda_{a_0})_{u_0}(\xi'(t))}.\end{aligned}\tag{3.3}$$

Since $\xi'(t) = (\widetilde{H}^{\widetilde{\xi(t)}^L})_{u_0+\xi(t)}$, we have

$$\begin{aligned}
&\lim_{t\to T-0}(1-(\lambda_{a_0})_{u_0}(\xi(t)))(\lambda_{a_0})_{u_0}(\xi'(t))\\
&=\lim_{t\to T-0}\sum_{a=1}^{\bar{r}}\left(\frac{(m_a^e+m_a^o)(1-(\lambda_{a_0})_{u_0}(\xi(t)))}{\tan(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(\xi(t))))}\right.\\
&\qquad\qquad\left.+\frac{(m_a^e-m_a^o)(1-(\lambda_{a_0})_{u_0}(\xi(t)))}{\sin(\frac{\pi}{b_a}(1-(\lambda_a)_{u_0}(\xi(t))))}\right)\frac{\pi}{2b_a}\langle(\mathbf{n}_a)_{u_0},(\mathbf{n}_{a_0})_{u_0}\rangle\\
&=\frac{1}{2}\lim_{t\to T-0}\left((m_{a_0}^e+m_{a_0}^o)\cos^2(\frac{\pi}{b_{a_0}}(1-(\lambda_{a_0})_{u_0}(\xi(t))))\right.\\
&\qquad\qquad\left.+(m_{a_0}^e-m_{a_0}^o)\frac{1}{\cos(\frac{\pi}{b_a}(1-(\lambda_{a_0})_{u_0}(\xi(t))))}\right)||(\mathbf{n}_{a_0})_{u_0}||^2\\
&=m_{a_0}^e||(\mathbf{n}_{a_0})_{u_0}||^2,
\end{aligned}$$

which together with (3.3) deduces

$$\lim_{t\to T-0}||\widetilde{A}_v^t||_\infty^2(T-t)=\frac{(\lambda_{a_0})_{u_0}(v)^2}{2m_{a_0}^e||(\mathbf{n}_{a_0})_{u_0}||^2} \tag{3.4}$$

and hence

$$\lim_{t\to T-0}\max_{v\in S^\perp_{u_0+\xi(t)}\widetilde{M'_t}}||\widetilde{A}_v^t||_\infty^2(T-t)=\frac{1}{2m_{a_0}^e}.$$

Thus the mean curvature flow $\widetilde{M'_t}$ has type I singularity. Set $\bar{v}_t := (\pi\circ\phi)_{*u_0+\xi(t)}(v)$ and let $\{\lambda_1^t,\cdots,\lambda_n^t\}$ ($\lambda_1^t\leq\cdots\leq\lambda_n^t$) (resp. $\{\mu_1^t,\cdots,\mu_n^t\}$ ($0\leq\mu_1^t\leq\cdots\leq\mu_n^t$)) be all the eigenvalues of $A^t_{\bar{v}_t}$ (resp. $R(\cdot,\bar{v}_t)\bar{v}_t$), where $n:=\dim M$. Since M is an irreducible equifocal submanifold of codimension greater than one by the assumption, it is homogeneous by the homogeneity theorem of Christ (see [Ch]) and hence it is a principal orbit of a Hermann action by the result of Heintze-Palais-Terng-Thorbergsson (see [HPTT]) and the classification of hyperpolar actions by Kollross (see [Kol]). Furthermore, M and its parallel submanifolds are curvature-adapted by the result of Goertsches-Thorbergsson (see [GT]). Therefore, $A^t_{\bar{v}_t}$ and $R(\cdot,\bar{v}_t)\bar{v}_t$ commute and hence we have

$$\sum_{i=1}^n\sum_{j=1}^n\left(\mathrm{Ker}(A^t_{\bar{v}_t}-\lambda_i^t\,\mathrm{id})\cap\mathrm{Ker}(R(\cdot,\bar{v}_t)\bar{v}_t-\mu_j^t\,\mathrm{id})\right)=T_{(\pi\circ\phi)(u_0+\xi(t))}M'_t.$$

Set $\bar{E}^t_{ij}:=\mathrm{Ker}(A^t_{\bar{v}_t}-\lambda_i^t\,\mathrm{id})\cap\mathrm{Ker}(R(\cdot,\bar{v}_t)\bar{v}_t-\mu_j^t\,\mathrm{id})$ ($i,j\in\{1,\cdots,n\}$) and $I_t :=$

$\{(i,j)\in\{1,\cdots,n\}^2\,|\,\bar{E}^t_{ij}\neq\{0\}\}$. For each $(i,j)\in I_t$, we have

$$\mathrm{Spec}(\widetilde{A}^t_v|_{(\pi\circ\phi)_*^{-1}(\bar{E}^t_{ij})})=\left\{\begin{array}{ll}\left\{\dfrac{\sqrt{\mu^t_j}}{\arctan\frac{\sqrt{\mu^t_j}}{\lambda^t_i}+k\pi}\,|\,k\in\mathbb{Z}\right\} & (\mu^t_j\neq 0)\\ \{\lambda^t_i\} & (\mu^t_j=0)\end{array}\right.$$

in terms of Proposition 3.2 of [Koi1] and hence

$$||\widetilde{A}^t_v||_\infty=\max\left(\left\{\frac{\sqrt{\mu^t_j}}{\arctan\frac{\sqrt{\mu^t_j}}{|\lambda^t_i|}}\,|\,(i,j)\in I_t\text{ s.t. }\mu^t_j\neq 0\right\}\cup\{|\lambda^t_i|\,|\,(i,j)\in I_t\text{ s.t. }\mu^t_j=0\}\right).$$

It is clear that $\sup\limits_{0\leq t<T}\mu^t_n<\infty$. If $\lim\limits_{t\to T-0}|\lambda^t_i|=\infty$, then we have $\lim\limits_{t\to T-0}\left(\dfrac{\sqrt{\mu^t_j}}{\arctan\frac{\sqrt{\mu^t_j}}{\lambda^t_i}}\right)/\lambda^t_i$ $=1$. Hence we have

$$\begin{aligned}&\lim_{t\to T-0}||\widetilde{A}^t_v||^2_\infty(T-t)\\ &=\max\left\{\lim_{t\to T-0}(\lambda^t_i)^2(T-t)\,|\,i=1,\cdots,n\right\}\\ &=\lim_{t\to T-0}\max\{(\lambda^t_i)^2(T-t)\,|\,i=1,\cdots,n\}\\ &=\lim_{t\to T-0}||A^t_{\bar{v}_t}||^2_\infty(T-t),\end{aligned}$$

which together with (3.4) deduces

$$\lim_{t\to T-0}||A^t_{\bar{v}_t}||^2_\infty(T-t)=\frac{(\lambda_{a_0})_{u_0}(v)^2}{2m^e_{a_0}||(\mathbf{n}_{a_0})_{u_0}||^2}.$$

Therefore we obtain

$$\lim_{t\to T-0}\max_{v\in S^\perp_{\exp^\perp(\xi(t))}M'_t}||A^t_v||^2_\infty(T-t)=\frac{1}{2m^e_{a_0}}<\infty.$$

Thus the mean curvature flow M'_t has type I singularity. q.e.d.

Next we prove Theorem B.

Proof of Theorem B. For simplicity, set $I:=\{1,\cdots,\bar{r}\}$, where $\bar{r}$ is as above. Let $\widetilde{\sigma}$ be a stratum of dimension greater than zero of $\partial\widetilde{C}$ and $I_{\widetilde{\sigma}}:=\{a\in I\,|\,\widetilde{\sigma}\subset(\lambda_a)^{-1}_{u_0}(1)\}$.

Let $w \in \widetilde{\sigma}$ and $\widetilde{w}$ be the parallel normal vector field of M with $\widetilde{w}_{x_0} = w$. The horizontal lift $\widetilde{w}^L$ of $\widetilde{w}$ to $H^0([0,1],\mathfrak{g})$ is a parallel normal vector field of $\widetilde{M}$. Denote by F (resp. $\widetilde{F}$) the focal submanifold of M (resp. $\widetilde{M}$) for $\widetilde{w}$ (resp. $\widetilde{w}^L$) and by $\widetilde{A}^{\widetilde{F}}$ (resp. $\widetilde{H}^{\widetilde{F}}$) the shape tensor (resp. the mean curvature vector) of $\widetilde{F}$. Then we have

$$T_{u_0+w}\widetilde{F} = \left(\bigoplus_{a\in I\setminus I_{\widetilde{\sigma}}} \bigoplus_{j\in\mathbf{Z}} \eta_{\widetilde{w}*}(E_{aj})\right) \oplus \left(\bigoplus_{a\in I_{\widetilde{\sigma}}} \bigoplus_{j\in\mathbf{Z}\setminus\{0\}} \eta_{\widetilde{w}*}(E_{aj})\right).$$

Also we have

$$T^{\perp}_{u_0+w}\widetilde{F} = \left(\bigoplus_{a\in I_{\widetilde{\sigma}}} (E_{a0})_{u_0}\right) \oplus T^{\perp}_{u_0}\widetilde{M},$$

where we identify $T_{u_0+w}H^0([0,1],\mathfrak{g})$ with $T_{u_0}H^0([0,1],\mathfrak{g})$. For $v \in T^{\perp}_{u_0}\widetilde{M}\,(\subset T^{\perp}_{u_0+w}\widetilde{F})$, we have

$$\widetilde{A}^{\widetilde{F}}_v|_{\eta_{\widetilde{w}*}((E_{aj})_{u_0})} = \frac{(\lambda_{aj})_{u_0}(v)}{1-(\lambda_{aj})_{u_0}(w)}\mathrm{id} \quad ((a,j) \in ((I \setminus I_{\widetilde{\sigma}}) \times \mathbb{Z}) \cup (I_{\widetilde{\sigma}} \times (\mathbb{Z} \setminus \{0\}))).$$

Hence we have

$$\begin{aligned}
&\mathrm{Tr}_r \widetilde{A}^{\widetilde{F}}_v \\
=& \sum_{a\in I\setminus I_{\widetilde{\sigma}}} \left(\sum_{j\in\mathbf{Z}} \frac{m^e_a(\lambda_{a,2j})_{u_0}(v)}{1-(\lambda_{a,2j})_{u_0}(w)} + \sum_{j\in\mathbf{Z}} \frac{m^o_a(\lambda_{a,2j+1})_{u_0}(v)}{1-(\lambda_{a,2j+1})_{u_0}(w)} \right) \\
&+ \sum_{a\in I_{\widetilde{\sigma}}} \left(\sum_{j\in\mathbf{Z}\setminus\{0\}} \frac{m^e_a(\lambda_{a,2j})_{u_0}(v)}{1-(\lambda_{a,2j})_{u_0}(w)} + \sum_{j\in\mathbf{Z}} \frac{m^o_a(\lambda_{a,2j+1})_{u_0}(v)}{1-(\lambda_{a,2j+1})_{u_0}(w)} \right) \\
=& \sum_{a\in I\setminus I_{\widetilde{\sigma}}} \Bigg((m^e_a + m^o_a)\cot(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a}) \\
&\qquad + (m^e_a - m^o_a)\mathrm{cosec}(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a}) \Bigg) \frac{\pi}{2b_a}(\lambda_a)_{u_0}(v) \\
&+ \sum_{a\in I_{\widetilde{\sigma}}} \left(\frac{m^e_a(\lambda_a)_{u_0}(v)}{b_a} \sum_{j\in\mathbf{Z}\setminus\{0\}} \frac{1}{2j} + \frac{m^o_a(\lambda_a)_{u_0}(v)}{b_a} \sum_{j\in\mathbf{Z}} \frac{1}{2j+1} \right) \\
=& \sum_{a\in I\setminus I_{\widetilde{\sigma}}} \Bigg((m^e_a + m^o_a)\cot(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a}) \\
&\qquad + (m^e_a - m^o_a)\mathrm{cosec}(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a}) \Bigg) \frac{\pi}{2b_a}(\lambda_a)_{u_0}(v),
\end{aligned}$$

that is, the $T^{\perp}_{u_0}\widetilde{M}$-component $((\widetilde{H}^{\widetilde{F}})_{u_0+w})_{T^{\perp}_{u_0}\widetilde{M}}$ of $(\widetilde{H}^{\widetilde{F}})_{u_0+w}$ is equal to

$$\begin{aligned}\sum_{a\in I\setminus I_{\widetilde{\sigma}}}\Bigg((m_a^e+m_a^o)\cot(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a})&\\ +(m_a^e-m_a^o)\mathrm{cosec}(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a})\Bigg)\frac{\pi}{2b_a}(\mathbf{n}_a)_{u_0}.&\end{aligned}$$

Denote by Φ_{u_0+w} the normal holonomy group of $\widetilde{F}$ at u_0+w and L_{u_0} be the focal leaf through u_0 for $\widetilde{w}$. Since $L_{u_0}=\Phi_{u_0+w}\cdot u_0$, there exists $\mu\in\Phi_{u_0+w}$ such that $\mu(T^{\perp}_{u_0}\widetilde{M})=T^{\perp}_{u_1}\widetilde{M}$ for any point u_1 of L_{u_0}. From this fact, we have $\mu(((\widetilde{H}^{\widetilde{F}})_{u_0+w})_{T^{\perp}_{u_0}\widetilde{M}})=((\widetilde{H}^{\widetilde{F}})_{u_0+w})_{T^{\perp}_{u_1}\widetilde{M}}$. Furthermore, from this fact, we have $(\widetilde{H}^{\widetilde{F}})_{u_0+w}\in\bigcap\limits_{u\in L_{u_0}}T^{\perp}_u\widetilde{M}$, which contains $\widetilde{\sigma}$ as an open subset. Therefore, we obtain

$$\begin{aligned}(\widetilde{H}^{\widetilde{F}})_{u_0+w}=\sum_{a\in I\setminus I_{\widetilde{\sigma}}}\Bigg((m_a^e+m_a^o)\cot(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a})&\\ +(m_a^e-m_a^o)\mathrm{cosec}(\frac{\pi(1-(\lambda_a)_{u_0}(w))}{b_a})\Bigg)\frac{\pi}{2b_a}(\mathbf{n}_a)_{u_0}\quad(\in T\widetilde{\sigma}).&\end{aligned}\tag{3.5}$$

Define a tangent vector field $X^{\widetilde{\sigma}}$ on $\widetilde{\sigma}$ by $X^{\widetilde{\sigma}}_w:=(\widetilde{H}^{\widetilde{F}})_{u_0+w}$ $(w\in\widetilde{\sigma})$. Fix $w\in\widetilde{\sigma}$. Let $\xi:[0,T)\to\widetilde{\sigma}$ be the (maximal) integral curve of $X^{\widetilde{\sigma}}$ with $\xi(0)=w$, where T is possible to be equal to ∞. Since $\xi(t)-w\in\widetilde{\sigma}\,(\subset T^{\perp}_{u_0+w}\widetilde{F})$, Φ_{u_0+w} preserves $\widetilde{\sigma}$ invariantly and it acts on $\widetilde{\sigma}$ trivially, $\xi(t)-w$ extends to a parallel normal vector field of $\widetilde{F}$. Denote by $\widetilde{\xi(t)-w}$ this parallel normal vector field and by $\widehat{\xi(t)-w}:=(\pi\circ\phi)_*(\widetilde{\xi(t)-w})$, which is a parallel normal vector field of F. Also, denote by $\widetilde{F}_{\widetilde{\xi(t)-w}}$ the parallel submanifold of $\widetilde{F}$ for $\widetilde{\xi(t)-w}$ and by $F_{\widehat{\xi(t)-w}}$ the parallel submanifold of F for $\widehat{\xi(t)-w}$. Let F_t (resp. $\widetilde{F}_t$) be the mean curvature flow having F (resp. $\widetilde{F}$) as initial data. Then, by imitating the proof of Lemma 3.1, we can show $\widetilde{F}_t=\widetilde{F}_{\widetilde{\xi(t)-w}}$ and $F_t=F_{\widehat{\xi(t)-w}}$ $(t\in[0,T))$. By using these facts and (3.5) and imitating the proof of the statement (i) of Theorem A, we can show the statement (i). Also, by imitating the proof of the statement (ii) of Theorem A, we can show the statement (ii). q.e.d.


$\widetilde{F}$
L_{u_0}
$\widetilde{M}$
$T^{\perp}_{u_0}\widetilde{M}$
u_0+w
$T^{\perp}_{u_1}\widetilde{M}$
$T^{\perp}_{u_0+w}\widetilde{F}$
u_0
u_1
$\bigcap_{u\in L_{u_0}} T^{\perp}_u\widetilde{M}$
$\widetilde{C}$
$(\widetilde{H}^F)_{u_0+w}$
$\widetilde{\sigma}$
u_0
$T^{\perp}_{u_1}\widetilde{M}$
$T^{\perp}_{u_0}\widetilde{M}$

Fig. 3.

4 The mean curvature flows of orbits of Hermann actions

In this section, we describe explicitly the mean curvature flows of orbits of Hermann actions whose principal orbits are full and irreducible. Let G/K be a symmetric space of compact type and H be a symmetric subgroup of G. Also, let θ be an involution of G with $(\mathrm{Fix}\,\theta)_0 \subset K \subset \mathrm{Fix}\,\theta$ and τ be an invloution of G with $(\mathrm{Fix}\,\tau)_0 \subset H \subset \mathrm{Fix}\,\tau$, where $\mathrm{Fix}\,\theta$ (resp. $\mathrm{Fix}\,\tau$) is the fixed point group of θ (resp. τ) and $(\mathrm{Fix}\,\theta)_0$ (resp. $(\mathrm{Fix}\,\tau)_0$) is the identity component of $\mathrm{Fix}\,\theta$ (resp. $\mathrm{Fix}\,\tau$). We assume that $\tau\circ\theta = \theta\circ\tau$. Set $L := \mathrm{Fix}(\theta\circ\tau)$. Denote by the same symbol θ (resp. τ) the involution of the Lie algebra $\mathfrak{g}$ of G induced from θ (resp. τ). Set $\mathfrak{k} := \mathrm{Ker}(\theta - \mathrm{id})$, $\mathfrak{p} := \mathrm{Ker}(\theta + \mathrm{id})$, $\mathfrak{h} := \mathrm{Ker}(\tau - \mathrm{id})$ and $\mathfrak{q} := \mathrm{Ker}(\tau + \mathrm{id})$. The space $\mathfrak{p}$ is identified with $T_{eK}(G/K)$. From $\theta\circ\tau = \tau\circ\theta$, we have $\mathfrak{p} = \mathfrak{p}\cap\mathfrak{h} + \mathfrak{p}\cap\mathfrak{q}$. Take a maximal abelian subspace $\mathfrak{b}$ of $\mathfrak{p}\cap\mathfrak{q}$ and let $\mathfrak{p} = \mathfrak{z}_{\mathfrak{p}}(\mathfrak{b}) + \sum\limits_{\beta\in\triangle'_+} \mathfrak{p}_\beta$ be the root space decomposition with respect to $\mathfrak{b}$, where $\mathfrak{z}_{\mathfrak{p}}(\mathfrak{b})$ is the centralizer of $\mathfrak{b}$ in $\mathfrak{p}$, $\triangle'_+$ is the positive root system of $\triangle' := \{\beta \in \mathfrak{b}^*\setminus\{0\} \,|\, \exists\, X(\neq 0) \in \mathfrak{p} \text{ s.t. } \mathrm{ad}(b)^2(X) = -\beta(b)^2 X \ (\forall\, b \in \mathfrak{b})\}$ under some lexicographic ordering of $\mathfrak{b}^*$ and $\mathfrak{p}_\beta := \{X \in \mathfrak{p} \,|\, \mathrm{ad}(b)^2(X) = -\beta(b)^2 X \ (\forall\, b\in\mathfrak{b})\}$ $(\beta \in \triangle'_+)$. Also, let

$\triangle'^V_+ := \{\beta \in \triangle'_+ \,|\, \mathfrak{p}_\beta \cap \mathfrak{q} \neq \{0\}\}$ and $\triangle'^H_+ := \{\beta \in \triangle'_+ \,|\, \mathfrak{p}_\beta \cap \mathfrak{h} \neq \{0\}\}$. Then we have $\mathfrak{q} = \mathfrak{b} + \sum\limits_{\beta \in \triangle'^V_+} (\mathfrak{p}_\beta \cap \mathfrak{q})$ and $\mathfrak{h} = \mathfrak{z}_{\mathfrak{h}}(\mathfrak{b}) + \sum\limits_{\beta \in \triangle'^H_+} (\mathfrak{p}_\beta \cap \mathfrak{h})$, where $\mathfrak{z}_{\mathfrak{h}}(\mathfrak{b})$ is the centralizer of $\mathfrak{b}$ in $\mathfrak{h}$. The orbit $H(eK)$ is a reflective submanifold and it is isometric to the symmetric space $H/H \cap K$ (equipped with a metric scaled suitably). Also, $\exp^\perp(T^\perp_{eK}(H(eK)))$ is also a reflective submanifold and it is isometric to the symmetric space $L/H \cap K$ (equipped with a metric scaled suitably), where $\exp^\perp$ is the normal exponential map of $H(eK)$. The system $\triangle'^V := \triangle'^V_+ \cup (-\triangle'^V_+)$ is the root system of $L/H \cap K$. Define a subset $\widetilde{C}$ of $\mathfrak{b}$ by

$$\begin{aligned} \widetilde{C} := \{b \in \mathfrak{b} \,|\, & 0 < \beta(b) < \pi \, (\forall \beta \in \triangle'^V_+ \setminus \triangle'^H_+),\ -\frac{\pi}{2} < \beta(b) < \frac{\pi}{2} \, (\forall \beta \in \triangle'^H_+ \setminus \triangle'^V_+), \\ & 0 < \beta(b) < \frac{\pi}{2} \, (\forall \beta \in \triangle'^V_+ \cap \triangle'^H_+)\}. \end{aligned}$$

Let Π be the simple root system of $\triangle'_+$, and set $\Pi_V := \Pi \cap \triangle'^V_+$ and $\Pi_H := \Pi \cap \triangle'^H_+$. Also, let δ be the highest root of $\triangle'^V_+ \cup 2\triangle'^H_+$. Set $\widetilde{\Pi} := \Pi \cup \{\delta\}$, $\widetilde{\Pi}_V := \widetilde{\Pi} \cap {\triangle'_+}^V$ and $\widetilde{\Pi}_H := \widetilde{\Pi} \cap ({\triangle'_+}^H \cup 2{\triangle'_+}^H)$. Then we have

$$\widetilde{C} = \{b \in \mathfrak{b} \,|\, \beta(b) > 0 \, (\forall \beta \in \Pi_V),\ \beta(b) > -\frac{\pi}{2} \, (\forall \beta \in \Pi_H \setminus \Pi_V),\ \delta(b) < \pi\},$$

Set $C := \mathrm{Exp}(\widetilde{C})$, where Exp is the exponential map of G/K at eK. Let $P(G, H \times K) := \{g \in H^1([0,1], G) \,|\, (g(0), g(1)) \in H \times K\}$, where $H^1([0,1], G)$ is the Hilbert Lie group of all H^1-paths in G. This group acts on $H^0([0,1], \mathfrak{g})$ as gauge action. The orbits of the $P(G, H \times K)$-action are the inverse images of orbits of the H-action by $\pi \circ \phi$. The set $\Sigma := \mathrm{Exp}(\mathfrak{b})$ is a section of the H-action and $\mathfrak{b}$ is a section of the $P(G, H \times K)$-action on $H^0([0,1], \mathfrak{g})$, where $\mathfrak{b}$ is identified with the horizontal lift of $\mathfrak{b}$ to the zero element $\hat{0}$ of $H^0([0,1], \mathfrak{g})$ ($\hat{0}$: the constant path at the zero element 0 of $\mathfrak{g}$). The set $\widetilde{C}$ is the fundamental domain of the Coxeter group of a principal $P(G, H \times K)$-orbit (hence a principal H-orbit) and each prinicipal H-orbit meets C at one point and each singular H-orbit meets ∂C at one point. The focal set of the principal orbit $P(G, H \times K) \cdot Z_0$ ($Z_0 \in \widetilde{C}$) at Z_0 consists of the hyperplanes $\beta^{-1}(j\pi)$'s ($\beta \in \triangle'^V_+ \setminus \triangle'^H_+,\ j \in \mathbb{Z}$), $\beta^{-1}((j + \frac{1}{2})\pi)$'s ($\beta \in \triangle'^H_+ \setminus \triangle'^V_+,\ j \in \mathbb{Z}$), $\beta^{-1}(\frac{j\pi}{2})$'s ($\beta \in \triangle'^V_+ \cap \triangle'^H_+,\ j \in \mathbb{Z}$) in $\mathfrak{b}(= T^\perp_{Z_0}(P(G, H \times K) \cdot Z_0))$. Denote by $\exp^G$ the exponential map of G. Note that $\pi \circ \exp^G|_{\mathfrak{p}} = \mathrm{Exp}$. Let $Y_0 \in \widetilde{C}$ and $M(Y_0) := H(\mathrm{Exp}(Y_0))$. Then we have $T^\perp_{\mathrm{Exp}(Y_0)} M(Y_0) = (\exp^G(Y_0))_*(\mathfrak{b})$. Denote by A^{Y_0} the shape tensor of $M(Y_0)$. Take $v \in T^\perp_{\mathrm{Exp}(Y_0)} M(Y_0)$ and set $\bar{v} := (\exp^G(Y_0))_*^{-1}(v)$. By scaling the metric of G/K by a suitable positive constant, we have

$$A^{Y_0}_v|_{\exp^G(Y_0)_*(\mathfrak{p}_\beta \cap \mathfrak{q})} = -\frac{\beta(\bar{v})}{\tan \beta(Y_0)} \mathrm{id} \quad (\beta \in \triangle'^V_+) \tag{4.1}$$

and

$$A^{Y_0}_v|_{\exp^G(Y_0)_*(\mathfrak{p}_\beta\cap\mathfrak{h})} = \beta(\bar{v})\tan\beta(Y_0)\mathrm{id}\quad(\beta\in\triangle'^H_+). \tag{4.2}$$

Set $m^V_\beta := \dim(\mathfrak{p}_\beta\cap\mathfrak{q})$ $(\beta\in\triangle'^V_+)$ and $m^H_\beta := \dim(\mathfrak{p}_\beta\cap\mathfrak{h})$ $(\beta\in\triangle'^H_+)$. Set $\widetilde{M}(Y_0) := (\pi\circ\phi)^{-1}(M(Y_0))(= P(G,H\times K)\cdot Y_0)$. We can show $(\pi\circ\phi)(Y_0) = \mathrm{Exp}(Y_0)$.



Fig. 4.

Denote by $\widetilde{A}^{Y_0}$ the shape tensor of $\widetilde{M}(Y_0)$. According to Proposition 3.2 of [Koi1], we have

$$\mathrm{Spec}(\widetilde{A}^{Y_0}_{\bar{v}}|_{(\pi\circ\phi)^{-1}_{*Y_0}(\exp^G(Y_0)_*(\mathfrak{p}_\beta\cap\mathfrak{q}))})\setminus\{0\} = \{\frac{-\beta(\bar{v})}{\beta(Y_0)+j\pi}\,|\,j\in\mathbb{Z}\}\quad(\beta\in\triangle'^V_+),$$
$$\mathrm{Spec}(\widetilde{A}^{Y_0}_{\bar{v}}|_{(\pi\circ\phi)^{-1}_{*Y_0}(\exp^G(Y_0)_*(\mathfrak{p}_\beta\cap\mathfrak{h}))})\setminus\{0\} = \{\frac{-\beta(\bar{v})}{\beta(Y_0)+(j+\frac{1}{2})\pi}\,|\,j\in\mathbb{Z}\}\quad(\beta\in\triangle'^H_+),$$

and

$$\mathrm{Spec}(\widetilde{A}^{Y_0}_{\bar{v}}|_{(\pi\circ\phi)^{-1}_{*Y_0}(\exp^G(Y_0)_*(\mathfrak{z}_{\mathfrak{h}}(\mathfrak{b})))}) = \{0\}.$$

Hence the set $\mathcal{PC}_{\widetilde{M}(Y_0)}$ of all principal curvatures of $\widetilde{M}(Y_0)$ is given by

$$\mathcal{PC}_{\widetilde{M}(Y_0)}=\{\frac{-\widetilde{\beta}}{\beta(Y_0)+j\pi}\,|\,\beta\in\triangle'^V_+,\ j\in\mathbb{Z}\}\cup\{\frac{-\widetilde{\beta}}{\beta(Y_0)+(j+\frac{1}{2})\pi}\,|\,\beta\in\triangle'^H_+,\ j\in\mathbb{Z}\},$$

where $\widetilde{\beta}$ is the parallel section of $(T^{\perp}\widetilde{M}(Y_0))^*$ with $\widetilde{\beta}_{u_0}=\beta\circ\exp^G(Y_0)_*^{-1}$. Also, we can show that the multiplicity of $\frac{-\widetilde{\beta}}{\beta(Y_0)+j\pi}$ $(\beta\in\triangle'^V_+)$ is equal to m^V_β and that of $\frac{-\widetilde{\beta}}{\beta(Y_0)+(j+\frac{1}{2})\pi}$ $(\beta\in\triangle'^H_+)$ is equal to m^H_β. Define $\lambda^{Y_0}_\beta$ and $b^{Y_0}_\beta$ $(\beta\in\triangle'_+)$ by

$$(\lambda^{Y_0}_\beta,b^{Y_0}_\beta):=\begin{cases}(\frac{-\widetilde{\beta}}{\beta(Y_0)},\frac{\pi}{\beta(Y_0)}) & (\beta\in\triangle'^V_+\setminus\triangle'^H_+)\\ (\frac{-\widetilde{\beta}}{\beta(Y_0)+\frac{\pi}{2}},\frac{\pi}{\beta(Y_0)+\frac{\pi}{2}}) & (\beta\in\triangle'^H_+\setminus\triangle'^V_+)\\ (\frac{-\widetilde{\beta}}{\beta(Y_0)},\frac{\pi}{2\beta(Y_0)}) & (\beta\in\triangle'^V_+\cap\triangle'^H_+).\end{cases}$$

Then we have $\frac{-\widetilde{\beta}}{\beta(Y_0)+j\pi}=\frac{\lambda^{Y_0}_\beta}{1+jb^{Y_0}_\beta}$ when $\beta\in\triangle'^V_+\setminus\triangle'^H_+$, $\frac{-\widetilde{\beta}}{\beta(Y_0)+(j+\frac{1}{2})\pi}=\frac{\lambda^{Y_0}_\beta}{1+jb^{Y_0}_\beta}$ when $\beta\in\triangle'^H_+\setminus\triangle'^V_+$ and $(\frac{-\widetilde{\beta}}{\beta(Y_0)+j\pi},\frac{-\widetilde{\beta}}{\beta(Y_0)+(j+\frac{1}{2})\pi})=(\frac{\lambda^{Y_0}_\beta}{1+2jb^{Y_0}_\beta},\frac{\lambda^{Y_0}_\beta}{1+(2j+1)b^{Y_0}_\beta})$ when $\beta\in\triangle'^V_+\cap\triangle'^H_+$. That is, we have

$$\mathcal{PC}_{\widetilde{M}(Y_0)}=\{\frac{\lambda^{Y_0}_\beta}{1+jb^{Y_0}_\beta}\,|\,\beta\in\triangle'_+,\ j\in\mathbb{Z}\}.$$

Denote by $m_{\beta j}$ the multiplicity of $\frac{\lambda_\beta}{1+jb_\beta}$. Then we have

$$m_{\beta,2j}=\begin{cases}m^V_\beta & (\beta\in\triangle'^V_+\setminus\triangle'^H_+)\\ m^H_\beta & (\beta\in\triangle'^H_+\setminus\triangle'^V_+)\\ m^V_\beta & (\beta\in\triangle'^V_+\cap\triangle'^H_+),\end{cases}\qquad m_{\beta,2j+1}=\begin{cases}m^V_\beta & (\beta\in\triangle'^V_+\setminus\triangle'^H_+)\\ m^H_\beta & (\beta\in\triangle'^H_+\setminus\triangle'^V_+)\\ m^H_\beta & (\beta\in\triangle'^V_+\cap\triangle'^H_+),\end{cases}$$

where $j\in\mathbb{Z}$. Denote by $\widetilde{H}^{Y_0}$ the mean curvature vector of $\widetilde{M}(Y_0)$ and $\mathbf{n}^{Y_0}_\beta$ the curvature normal corresponding to $\lambda^{Y_0}_\beta$. From (3.1) (the case of $v=0$), we have

$$\begin{aligned}
\widetilde{H}^{Y_0} = & \sum_{\beta\in\triangle'^V_+\setminus\triangle'^H_+} 2m^V_\beta\cot\frac{\pi}{b^{Y_0}_\beta}\times\frac{\pi}{2b^{Y_0}_\beta}\mathbf{n}^{Y_0}_\beta \\
& + \sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+} 2m^H_\beta\cot\frac{\pi}{b^{Y_0}_\beta}\times\frac{\pi}{2b^{Y_0}_\beta}\mathbf{n}^{Y_0}_\beta \\
& + \sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+}\left((m^V_\beta+m^H_\beta)\cot\frac{\pi}{b^{Y_0}_\beta}+(m^V_\beta-m^H_\beta)\mathrm{cosec}\frac{\pi}{b^{Y_0}_\beta}\right)\frac{\pi}{2b^{Y_0}_\beta}\mathbf{n}^{Y_0}_\beta \\
= & \sum_{\beta\in\triangle'^V_+\setminus\triangle'^H_+} m^V_\beta\beta(Y_0)\cot\beta(Y_0)\mathbf{n}^{Y_0}_\beta \\
& - \sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+} m^H_\beta(\beta(Y_0)+\frac{\pi}{2})\tan\beta(Y_0)\mathbf{n}^{Y_0}_\beta \\
& + \sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+}\beta(Y_0)\big((m^V_\beta+m^H_\beta)\cot 2\beta(Y_0) \\
& \qquad +(m^V_\beta-m^H_\beta)\mathrm{cosec}2\beta(Y_0)\big)\,\mathbf{n}^{Y_0}_\beta.
\end{aligned}\tag{4.3}$$

Define $\beta^\sharp\,(\in\mathfrak{b})$ by $\beta(\cdot)=\langle\beta^\sharp,\cdot\rangle$ and let $\widetilde{\beta^\sharp}^{Y_0}$ be the parallel normal vector field of $\widetilde{M}(Y_0)$ with $(\widetilde{\beta^\sharp}^{Y_0})_{Y_0}=\beta^\sharp$, where we identify $\mathfrak{b}$ with $T^\perp_{Y_0}\widetilde{M}(Y_0)$. From the definition of $\lambda^{Y_0}_\beta$, we have

$$\mathbf{n}^{Y_0}_\beta=\begin{cases} -\dfrac{1}{\beta(Y_0)}\widetilde{\beta^\sharp}^{Y_0} & (\beta\in\triangle'^V_+\setminus\triangle'^H_+\text{ or }\beta\in\triangle'^V_+\cap\triangle'^H_+) \\ -\dfrac{1}{\beta(Y_0)+\frac{\pi}{2}}\widetilde{\beta^\sharp}^{Y_0} & (\beta\in\triangle'^H_+\setminus\triangle'^V_+). \end{cases}$$

Substituting this relation into (4.3), we have

$$\begin{aligned}
\widetilde{H}^{Y_0} = & -\sum_{\beta\in\triangle'^V_+\setminus\triangle'^H_+} m^V_\beta\cot\beta(Y_0)\widetilde{\beta^\sharp}^{Y_0}+\sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+} m^H_\beta\tan\beta(Y_0)\widetilde{\beta^\sharp}^{Y_0} \\
& -\sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+}\big((m^V_\beta+m^H_\beta)\cot 2\beta(Y_0) \\
& \qquad +(m^V_\beta-m^H_\beta)\mathrm{cosec}2\beta(Y_0)\big)\,\widetilde{\beta^\sharp}^{Y_0}.
\end{aligned}\tag{4.4}$$

Define a tangent vector field X on $\widetilde{C}$ by $X_{Y_0}:=(\widetilde{H}^{Y_0})_{Y_0}\,(\in T^\perp_{Y_0}\widetilde{M}(Y_0)=\mathfrak{b}(\subset V))$.

From (4.4), we have

$$
\begin{aligned}
X_{Y_0} = & -\sum_{\beta\in\triangle'^V_+\setminus\triangle'^{\,H}_+} m^V_\beta \cot\beta(Y_0)\beta^\sharp + \sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+} m^H_\beta \tan\beta(Y_0)\beta^\sharp \\
& -\sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+} \big((m^V_\beta + m^H_\beta)\cot 2\beta(Y_0) \\
& \qquad\qquad +(m^V_\beta - m^H_\beta)\mathrm{cosec}2\beta(Y_0)\big)\,\beta^\sharp.
\end{aligned} \tag{4.5}
$$

In order to analyze the mean curvature flows having principal orbits of the H-actions as initial data, we have only to analyze this vector field X. By using (4.5), we can explicitly describe X for each Hermann action. Set

$$
l_\beta := \begin{cases} \beta^{-1}(0) & (\beta\in\Pi_V) \\ \beta^{-1}(-\frac{\pi}{2}) & (\beta\in\Pi_H\setminus\Pi_V) \\ \delta^{-1}(\pi) & (\beta=\delta). \end{cases}
$$

Let $a=1$ when $\delta\in\triangle'^V_+\setminus\triangle'^H_+$ and $a=\frac{1}{2}$ when $\delta\in\triangle'^H_+$. Fix a stratum $\widetilde{\sigma}$ of $\partial\widetilde{C}$. Define $\triangle'^{\widetilde{\sigma}}_+$ by $\triangle'^{\widetilde{\sigma}}_+ := (\triangle'_+\setminus(\Pi\cup\{a\delta\}))\cup\{\beta\in\Pi\cup\{a\delta\}\,|\,\widetilde{\sigma}\not\subset l_\beta\}$ and set $(\triangle'^V_+)^{\widetilde{\sigma}} := \triangle'^V_+\cap\triangle'^{\widetilde{\sigma}}_+$ and $(\triangle'^H_+)^{\widetilde{\sigma}} := \triangle'^H_+\cap\triangle'^{\widetilde{\sigma}}_+$. Fix $Z_0\in\widetilde{\sigma}$. Set $F(Z_0) := H(\mathrm{Exp}Z_0)$ and $\widetilde{F}(Z_0) := P(G, H\times K)\cdot Z_0$. Denote by $\widetilde{H}^{Z_0}$ (resp. H^{Z_0}) the mean curvature vector of $\widetilde{F}(Z_0)$ (resp. $F(Z_0)$). Define a tangent vector field $X^{\widetilde{\sigma}}$ on $\widetilde{\sigma}$ by $(X^{\widetilde{\sigma}})_{Z_0} := (\widetilde{H}^{Z_0})_{Z_0}$ ($Z_0\in\widetilde{\sigma}$). From (3.5) (the case of $w=Z_0-Y_0$), we have

$$
\begin{aligned}
(X^{\widetilde{\sigma}})_{Z_0} = & -\sum_{\beta\in(\triangle'^V_+)^{\widetilde{\sigma}}\setminus(\triangle'^H_+)^{\widetilde{\sigma}}} m^V_\beta\cot\beta(Z_0)\beta^\sharp \\
& +\sum_{\beta\in(\triangle'^H_+)^{\widetilde{\sigma}}\setminus(\triangle'^V_+)^{\widetilde{\sigma}}} m^H_\beta\tan\beta(Z_0)\beta^\sharp \\
& -\sum_{\beta\in(\triangle'^V_+)^{\widetilde{\sigma}}\cap(\triangle'^H_+)^{\widetilde{\sigma}}} \big((m^V_\beta+m^H_\beta)\cot 2\beta(Z_0) \\
& \qquad\qquad +(m^V_\beta-m^H_\beta)\mathrm{cosec}2\beta(Z_0)\big)\,\beta^\sharp.
\end{aligned} \tag{4.6}
$$

In order to analyze the mean curvature flows having singular orbits through $\widetilde{\sigma}$ of the H-action as initial data, we have only to analyze the vector field $X^{\widetilde{\sigma}}$. Now we shall describe explicitly the above vector fields X and $X^{\widetilde{\sigma}}$ for some Hermann actions. We shall use the above notations.

Example 1. We consider the isotropy action $Sp(n)\curvearrowright SU(2n)/Sp(n)$. Then, since $\triangle'$ is equal to the root system of $SU(2n)/Sp(n)$, it is of (A_{n-1})-type and each of its roots is of multiplicity 4. We can describe $\mathfrak{b}, \triangle'_+, \Pi$ and δ as

$$
\begin{aligned}
&\mathfrak{b} = \{\sum_{i=1}^n x_i e_i \,|\, \sum_{i=1}^n x_i = 0\} (\subset \mathbf{R}^n),\\
&\triangle'_+ = \{(\beta_i - \beta_j)|_{\mathfrak{b}} \,|\, 1 \le i < j \le n\},\\
&\Pi = \{(\beta_i - \beta_{i+1})|_{\mathfrak{b}} \,|\, 1 \le i \le n-1\}, \quad \delta = (\beta_1 - \beta_n)|_{\mathfrak{b}},
\end{aligned}
\tag{4.7}
$$

where $(e_1, \cdots, e_n)$ is the orthonormal base of $\mathbf{R}^n$ and $(\beta_1, \cdots, \beta_n)$ is the dual base of $(e_1, \cdots, e_n)$. For simplicity, we set $\beta_{ij} := (\beta_i - \beta_j)|_{\mathfrak{b}}\,(1 \le i < j \le n)$, $I := \{1, \cdots, n\}$ and $\widehat{I} := \{(i,j) \in I^2 \,|\, 1 \le i < j \le n\}$. Since we consider the isotropy action, we have $\triangle'^V_+ = \triangle'_+$ and $\triangle'^H_+ = \emptyset$, that is,

$$
\begin{aligned}
\widetilde{C} &= \{\mathbf{x} \in \mathfrak{b} \,|\, 0 < \beta_{ij}(\mathbf{x}) < \pi \ ((i,j) \in \widehat{I})\}\\
&= \{\mathbf{x} \in \mathfrak{b} \,|\, \beta_{i,i+1}(\mathbf{x}) > 0\,(i = 1, \cdots, n-1),\ \beta_{1n}(\mathbf{x}) < \pi\}.
\end{aligned}
$$

From (4.5), we can describe X explicitly as

$$
\begin{aligned}
X_{\mathbf{x}} &= -4 \sum_{(i,j)\in\widehat{I}} \cot \beta_{ij}(\mathbf{x}) \beta_{ij}^{\sharp} = -4 \sum_{(i,j)\in\widehat{I}} \cot(x_i - x_j)(e_i - e_j)\\
&= -4 \sum_{i \in I} (\sum_{j \in I \setminus \{i\}} \cot(x_i - x_j)) e_i \qquad (\mathbf{x}(= \sum_{i=1}^n x_i e_i) \in \widetilde{C}).
\end{aligned}
\tag{4.8}
$$

Take a stratum $\widetilde{\sigma}$ of $\partial\widetilde{C}$. Set $\widehat{I}_{\widetilde{\Pi}} := \{(i, i+1) \,|\, 1 \le i \le n-1\} \cup \{(1,n)\}$, $l_{i,i+1} := \beta_{i,i+1}^{-1}(0)$ $(1 \le i \le n-1)$ and $l_{1n} := \beta_{1n}^{-1}(\pi)$. Set $\widehat{I}_{\widetilde{\sigma}} := \{(i,j) \in \widehat{I}_{\widetilde{\Pi}} \,|\, \widetilde{\sigma} \subset l_{ij}\}$. Also, set $I_i^{\widetilde{\sigma}} := \{j \in I \,|\, (i,j) \in \widehat{I} \setminus \widehat{I}_{\widetilde{\sigma}} \text{ or } (j,i) \in \widehat{I} \setminus \widehat{I}_{\widetilde{\sigma}}\}$ $(i = 1, \cdots, n)$. Then, from (4.6), we can describe $X^{\widetilde{\sigma}}$ explicitly as

$$
\begin{aligned}
X_{\mathbf{x}}^{\widetilde{\sigma}} &= -4 \sum_{(i,j)\in\widehat{I}\setminus\widehat{I}_{\widetilde{\sigma}}} \cot(x_i - x_j)(e_i - e_j)\\
&= -4 \sum_{i \in I} \left(\sum_{j \in I_i^{\widetilde{\sigma}}} \cot(x_i - x_j) \right) e_i \qquad (\mathbf{x}(= \sum_{i=1}^n x_i e_i) \in \widetilde{\sigma}).
\end{aligned}
\tag{4.9}
$$

According to (4.8), a principal orbit $Sp(n)(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x} \in \widetilde{C})$ is minimal if and only if the following relations hold:

$$
\sum_{j \in I \setminus \{i\}} \cot(x_i - x_j) = 0 \quad (i = 1, \cdots, n).
\tag{4.10}
$$

Also, according to (4.9), a singular orbit $Sp(n)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{\sigma}$) is minimal if and only if the following relations hold:

$$\sum_{j \in I_i^{\widetilde{\sigma}}} \cot(x_i - x_j) = 0 \quad (i = 1, \cdots, n). \tag{4.11}$$

In the sequel, we consider the case of $n = 3$. By solving (4.10) under $0 < x_i - x_j < \pi$ ($(i,j) \in \widehat{I}$), we have $(x_1, x_2, x_3) = (\frac{\pi}{3}, 0, -\frac{\pi}{3})$. Therefore the orbit $Sp(3)(\mathrm{Exp}(\frac{\pi}{3}e_1 - \frac{\pi}{3}e_3))$ is the only minimal principal orbit of the $Sp(3)$-action. Denote by $\widetilde{\sigma}_{12}$, $\widetilde{\sigma}_{23}$ and $\widetilde{\sigma}_{13}$ one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $\beta_{23}^{-1}(0)$ and $\beta_{13}^{-1}(\pi)$, respectively. For $(k,l) \in \widehat{I}$, by solving (4.11) (the case of $\widetilde{\sigma} = \widetilde{\sigma}_{kl}$) under $0 < x_i - x_j < \pi$ ($(i,j) \in \widehat{I} \setminus \widehat{I}_{\widetilde{\sigma}_{kl}}$), we have

$$(x_1, x_2, x_3) = \begin{cases} (\frac{\pi}{6}, \frac{\pi}{6}, -\frac{\pi}{3}) & ((k,l) = (1,2)) \\ (\frac{\pi}{3}, -\frac{\pi}{6}, -\frac{\pi}{6}) & ((k,l) = (2,3)) \\ (\frac{\pi}{2}, 0, -\frac{\pi}{2}) & ((k,l) = (1,3)). \end{cases}$$

Therefore the orbits $Sp(3)(\mathrm{Exp}(\frac{\pi}{6}(e_1 + e_2 - 2e_3)))$, $Sp(3)(\mathrm{Exp}(\frac{\pi}{6}(2e_1 - e_2 - e_3)))$ and $Sp(3)(\mathrm{Exp}(\frac{\pi}{2}(e_1 - e_3)))$ are the only minimal singular orbits through one dimensional stratums of $\partial\widetilde{C}$ of the $Sp(3)$-action.


$\beta_{12}^{-1}(\pi)$
$\frac{\pi}{6}(2e_1 - e_2 - e_3)$
$\frac{\pi}{3}(e_1 - e_3)$
$\frac{\pi}{2}(e_1 - e_3)$
$\beta_{23}^{-1}(0)$
$\beta_{13}^{-1}(\pi)$
$\beta_{23}^{-1}(\pi)$
$\mathbf{0}$
$\frac{\pi}{6}(e_1 + e_2 - 2e_3)$
$\beta_{12}^{-1}(0)$

Fig. 5

Next we shall investigate the divergences $\mathrm{div}\,X$ and $\mathrm{div}\,X^{\widetilde{\sigma}_{kl}}$ of X and $X^{\widetilde{\sigma}_{kl}}$ ($(k,l) \in \widehat{I}$). Take an orthonormal base $\{v_1 := \frac{1}{\sqrt{2}}(e_1 - e_2),\ v_2 := \frac{1}{\sqrt{6}}(e_1 + e_2 - 2e_3)\}$ of $\mathfrak{b}$. Let

$\mathbf{x} = \sum_{i=1}^{3} x_i e_i = \sum_{i=1}^{2} y_i v_i$. Then, from (4.8), we have

$$\begin{aligned} X_{\mathbf{x}} = -4 & \left(\sqrt{2}\cot\sqrt{2}y_1 + (\sqrt{2} - \frac{1}{\sqrt{2}})\cot(\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2) \right. \\ & \left. -\frac{1}{\sqrt{2}}\cot(-\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2) \right) v_1 \\ & -2\sqrt{6}\left(\cot(\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2) + \cot(-\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2) \right) v_2 \end{aligned}$$

and hence

$$(\operatorname{div} X)_{\mathbf{x}} = 8\left(\frac{1}{\sin^2\sqrt{2}y_1} + \frac{1}{\sin^2(\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2)} + \frac{1}{\sin^2(-\frac{1}{\sqrt{2}}y_1 + \frac{\sqrt{6}}{2}y_2)} \right) > 0.$$

Therefore, since X has the only zero point, $\operatorname{div} X > 0$ on $\widetilde{C}$ and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_{kl}\,((k,l) \in \widehat{I})$ (see the proof of Theorem A), all the integral curves of X through points other than its zero point $\frac{\pi}{3}(e_1 - e_3)$ converge to points of $\partial\widetilde{C}$ in finite time. Similarly we can show $\operatorname{div} X^{\widetilde{\sigma}_{kl}} > 0$ on $\widetilde{\sigma}_{kl}$ $((k,l) \in \widehat{I})$. Hence all integral curves of $X^{\widetilde{\sigma}_{kl}}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_{kl}$ in finite time. Therefore we obtain the following fact.

Proposition 4.1. *The mean curvature flow having any non-minimal principal orbit of the* $Sp(3)$*-action (on* $SU(6)/Sp(3)$*) as initial data converges to some singular orbit in finite time. Also, the mean curvature flow having any non-minimal singular orbit through* $\exp^{\perp}(\widetilde{\sigma}_{kl})$ *of the* $Sp(3)$*-action as initial data converges to one of two singular orbits through* $\exp^{\perp}(\partial\widetilde{\sigma}_{kl})$ *in finite time.*

Example 2. We consider the Hermann action $SO(2n) \curvearrowright SU(2n)/Sp(n)$. Since $L/H \cap K$ is equal to $(SU(n) \times SU(n))/SU(n)$, the cohomogeneity of this action is equal to the rank $n-1$ of $(SU(n) \times SU(n))/SU(n)$. On the other hand, the rank of $SU(2n)/Sp(n)$ also is equal to $n-1$. Thus the cohomogeneity of this action is equal to the rank of $SU(2n)/Sp(n)$. Hence a maximal abelian subspace $\mathfrak{b}$ of $\mathfrak{p} \cap \mathfrak{q}$ is also a maximal abelian subspace of $\mathfrak{p}$ and hence $\triangle'$ is the root system of $SU(2n)/Sp(n)$, it is of (A_{n-1})-type and each of its roots is of multiplicity 4. Hence the quantities $\mathfrak{b}$, $\triangle'_+$, Π and δ are given as in (4.7). Let β_i, β_{ij}, I and $\widehat{I}$ be as in Example 1. Also, since $\triangle'^V$ is the root system of $(SU(n) \times SU(n))/SU(n)$, it is of (A_{n-1})-type and each of its roots is of multiplicity 2. Hence we have $\triangle'^H_+ = \triangle'_+$

and each of its roots is of multiplicity 2. Therefore we have

$$\widetilde{C} = \{\mathbf{x} \in \mathfrak{b} \,|\, \beta_{i,i+1}(\mathbf{x}) > 0 \ (i = 1, \cdots, n-1),\ \beta_{1n}(\mathbf{x}) < \frac{\pi}{2}\}$$

and

$$\begin{aligned} X_{\mathbf{x}} &= -4 \sum_{(i,j)\in\widehat{I}} \cot 2\beta_{ij}(\mathbf{x})\beta_{ij}^{\sharp} = -4 \sum_{(i,j)\in\widehat{I}} \cot 2(x_i - x_j)(e_i - e_j) \\ &= -4 \sum_{i\in I}\Big(\sum_{j\in I\setminus\{i\}} \cot 2(x_i - x_j)\Big)e_i. \end{aligned} \tag{4.12}$$

Take a stratum $\widetilde{\sigma}$ of $\partial\widetilde{C}$. Let $\widehat{I}_{\widetilde{\sigma}}$ and $I_i^{\widetilde{\sigma}}$ be as in Example 1, where $l_{1n} := \beta_{1n}^{-1}(\frac{\pi}{2})$. Then, from (4.6), we can describe $X^{\widetilde{\sigma}}$ explicitly as

$$X_{\mathbf{x}}^{\widetilde{\sigma}} = -4 \sum_{i\in I} \left(\sum_{j\in I_i^{\widetilde{\sigma}}} \cot 2(x_i - x_j) \right) e_i \qquad (\mathbf{x}(= \sum_{i=1}^{n} x_i e_i) \in \widetilde{\sigma}). \tag{4.13}$$

According to (4.12), a principal orbit $SO(2n)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{C}$) is minimal if and only if the following relations hold:

$$\sum_{j\in I\setminus\{i\}} \cot 2(x_i - x_j) = 0 \quad (i = 1, \cdots, n). \tag{4.14}$$

Also, according to (4.13), a singular orbit $Sp(n)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{\sigma}$) is minimal if and only if the following relations hold:

$$\sum_{j\in I_i^{\widetilde{\sigma}}} \cot 2(x_i - x_j) = 0 \quad (i = 1, \cdots, n). \tag{4.15}$$

In the sequel, we consider the case of $n = 3$. By solving (4.14) under $0 < x_i - x_j < \frac{\pi}{2}$ ($(i,j) \in \widehat{I}$), we have $(x_1, x_2, x_3) = (\frac{\pi}{6}, 0, -\frac{\pi}{6})$. Therefore the orbit $SO(6)(\mathrm{Exp}(\frac{\pi}{6}e_1 - \frac{\pi}{6}e_3))$ is the only minimal principal orbit of the $SO(6)$-action. Denote by $\widetilde{\sigma}_{12}$, $\widetilde{\sigma}_{23}$ and $\widetilde{\sigma}_{13}$ one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $\beta_{23}^{-1}(0)$ and $\beta_{13}^{-1}(\frac{\pi}{2})$, respectively. For $(k,l) \in \widehat{I}$, by solving (4.15) (the case of $\widetilde{\sigma} = \widetilde{\sigma}_{kl}$) under $0 < x_i - x_j < \frac{\pi}{2}$ ($(i,j) \in \widehat{I} \setminus \widehat{I}_{\widetilde{\sigma}_{kl}}$), we have

$$(x_1, x_2, x_3) = \begin{cases} (\frac{\pi}{12}, \frac{\pi}{12}, -\frac{\pi}{6}) & ((k,l) = (1,2)) \\ (\frac{\pi}{6}, -\frac{\pi}{12}, -\frac{\pi}{12}) & ((k,l) = (2,3)) \\ (\frac{\pi}{4}, 0, -\frac{\pi}{4}) & ((k,l) = (1,3)). \end{cases}$$

Therefore the orbits $SO(6)(\mathrm{Exp}(\frac{\pi}{12}(e_1+e_2-2e_3)))$, $SO(6)(\mathrm{Exp}(\frac{\pi}{12}(2e_1-e_2-e_3)))$ and $SO(6)(\mathrm{Exp}(\frac{\pi}{4}(e_1-e_3)))$ are the only minimal singular orbits through one dimensional stratums of $\partial\widetilde{C}$ of the $SO(6)$-action.


$\beta_{12}^{-1}(\frac{\pi}{2})$
$\frac{\pi}{12}(2e_1-e_2-e_3)$
$\frac{\pi}{6}(e_1-e_3)$
$\frac{\pi}{4}(e_1-e_3)$
$\beta_{23}^{-1}(0)$
$\beta_{13}^{-1}(\frac{\pi}{2})$
$\beta_{23}^{-1}(\frac{\pi}{2})$
$\mathbf{0}$
$\frac{\pi}{12}(e_1+e_2-2e_3)$
$\beta_{12}^{-1}(0)$

Fig. 6

Next we shall investigate the divergences $\operatorname{div} X$ and $\operatorname{div} X^{\widetilde{\sigma}_{kl}}$ $((k,l)\in\widehat{I})$. Take an orthonormal base $\{v_1,\, v_2\}$ of $\mathfrak{b}$ as in Example 1. Let $\mathbf{x}=\sum_{i=1}^{3}x_ie_i=\sum_{i=1}^{2}y_iv_i$. Then, from (4.12), we have

$$\begin{aligned}X_{\mathbf{x}}=-4\Big(&\sqrt{2}\cot 2\sqrt{2}y_1+(\sqrt{2}-\frac{1}{\sqrt{2}})\cot(\sqrt{2}y_1+\sqrt{6}y_2)\\&-\frac{1}{\sqrt{2}}\cot(-\sqrt{2}y_1+\sqrt{6}y_2)\Big)v_1\\&-2\sqrt{6}\Big(\cot(\sqrt{2}y_1+\sqrt{6}y_2)+\cot(-\sqrt{2}y_1+\sqrt{6}y_2)\Big)v_2\end{aligned}$$

and hence

$$(\operatorname{div} X)_{\mathbf{x}}=16\left(\frac{1}{\sin^2 2\sqrt{2}y_1}+\frac{1}{\sin^2(\sqrt{2}y_1+\sqrt{6}y_2)}+\frac{1}{\sin^2(-\sqrt{2}y_1+\sqrt{6}y_2)}\right)>0.$$

Therefore, since X has the only zero point and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_{kl}\,((k,l)\in\widehat{I})$ by the proof of Theorem A, all the integral curves of X through points other than its zero point $\frac{\pi}{6}(e_1-e_3)$ converge to points of $\partial\widetilde{C}$ in finite time. Similarly we can show $\operatorname{div} X^{\widetilde{\sigma}_{kl}}>0$ on $\widetilde{\sigma}_{kl}$ $((k,l)\in\widehat{I})$. Hence all integral curves of $X^{\widetilde{\sigma}_{kl}}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_{kl}$ in finite time. Therefore we obtain the following fact.

Proposition 4.2. *The mean curvature flow having any non-minimal principal orbit of the* $SO(6)$*-action (on* $SU(6)/Sp(3)$*) as initial data converges to some singular orbit in finite time. Also, the mean curvature flow having any non-minimal singular orbit through* $\exp^{\perp}(\widetilde{\sigma}_{kl})$ *of the* $SO(6)$*-action as initial data converges to one of two singular orbits through* $\exp^{\perp}(\partial\widetilde{\sigma}_{kl})$ *in finite time.*

Example 3. We consider the isotropy action $S(U(p)\times U(q)) \curvearrowright SU(p+q)/S(U(p)\times U(q))$ $(p<q)$. Then, since $\triangle'$ is the root system of $SU(p+q)/S(U(p)\times U(q))$, it is of (B_p)-type and $\mathfrak{b}, \triangle'_+, \Pi$ and δ are described as

$$
\begin{aligned}
&\mathfrak{b} = \mathrm{Span}\{e_1,\cdots,e_p\},\\
&\triangle'_+ = \{\beta_i-\beta_j \,|\, 1\le i<j\le p\}\cup\{\beta_i\,|\,1\le i\le p\}\\
&\qquad\quad \cup\{\beta_i+\beta_j\,|\,1\le i<j\le p\}\cup\{2\beta_i\,|\,1\le i\le p\},\\
&\Pi = \{\beta_i-\beta_{i+1}\,|\,1\le i\le p-1\}\cup\{\beta_p\},\ \delta=\beta_1+\beta_2,
\end{aligned}
\tag{4.16}
$$

where $(e_1,\cdots,e_p)$ is the orthonormal base of $\mathfrak{b}$ and $(\beta_1,\cdots,\beta_p)$ is the dual base of $(e_1,\cdots,e_p)$. For simplicity, we set $\beta_{ij} := \beta_i-\beta_j$ $(1\le i<j\le p)$, $I:=\{1,\cdots,p\}$ and $\widehat{I} := \{(i,j)\in I^2\,|\,1\le i<j\le p\}$. It is shown that β_{ij}'s $(1\le i<j\le p)$ and $\beta_i+\beta_j$'s $(1\le i<j\le p)$ are of multiplicity 2, β_i's $(1\le i\le p)$ are of multiplicity $2(q-p)$ and $2\beta_i$'s $(1\le i\le p)$ are of multiplicity 1. Since we consider the isotropy action, we have $\triangle'^V_+ = \triangle'_+$ and $\triangle'^H_+ = \emptyset$, that is,

$$
\begin{aligned}
\widetilde{C} &= \{\mathbf{x}\in\mathfrak{b}\,|\,0<\beta_{ij}(\mathbf{x})<\pi\ ((i,j)\in\widehat{I}),\ 0<(\beta_i+\beta_j)(\mathbf{x})<\pi\ ((i,j)\in\widehat{I}),\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad 0<2\beta_i(\mathbf{x})<\pi\ (i\in I)\}\\
&= \{\mathbf{x}\in\mathfrak{b}\,|\,\beta_{i,i+1}(\mathbf{x})>0\,(1\le i\le p-1),\,\beta_p(\mathbf{x})>0,\ (\beta_1+\beta_2)(\mathbf{x})<\pi\}.
\end{aligned}
$$

From (4.5), we can describe X explicitly as

$$
\begin{aligned}
X_{\mathbf{x}} &= -\sum_{(i,j)\in\widehat{I}} 2\cot(x_i-x_j)(e_i-e_j) - \sum_{(i,j)\in\widehat{I}} 2\cot(x_i+x_j)(e_i+e_j)\\
&\quad -\sum_{i\in I} 2(q-p)\cot x_i\cdot e_i - \sum_{i\in I}\cot 2x_i\cdot 2e_i\\
&= -2\sum_{i\in I}\left(\sum_{j\in I\setminus\{i\}}(\cot(x_i-x_j)) + \cot(x_i+x_j)) + (q-p)\cot x_i + \cot 2x_i\right)e_i\\
&\qquad\qquad\qquad\qquad\qquad (\mathbf{x}(=\sum_{i=1}^{n} x_i e_i)\in\widetilde{C}).
\end{aligned}
$$

According to this relation, a principal orbit $S(U(p)\times U(q))(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x}\in\widetilde{C}$) is minimal if and only if the following relations hold:

$$\sum_{j\in I\setminus\{i\}}(\cot(x_i-x_j)+\cot(x_i+x_j))+(q-p)\cot x_i+\cot 2x_i=0\quad(i=1,\cdots,p). \tag{4.17}$$

In the sequel, we consider the case of $p=2$. Then (4.17) is as follows:

$$\left\{\begin{array}{l}\cot(x_1-x_2)+\cot(x_1+x_2)+(q-2)\cot x_1+\cot 2x_1=0\\ \cot(x_2-x_1)+\cot(x_1+x_2)+(q-2)\cot x_2+\cot 2x_2=0.\end{array}\right. \tag{4.18}$$

Under $0<x_i-x_j<\pi$ ($(i,j)\in\widehat{I}$), $0<x_i+x_j<\pi$ ($(i,j)\in\widehat{I}$) and $0<x_i<\frac{\pi}{2}$ ($i\in I$), the equation (4.18) has the only solution, which we denote by (α_1,α_2). Therefore the orbit $S(U(2)\times U(q))(\mathrm{Exp}(\alpha_1e_1+\alpha_2e_2))$ is the only minimal principal orbit of the $S(U(2)\times U(q))$-action. Here we note that $\lim\limits_{q\to\infty}(\alpha_1,\alpha_2)=(\frac{\pi}{2},\frac{\pi}{2})$. Denote by $\widetilde{\sigma}_1$, $\widetilde{\sigma}_2$ and $\widetilde{\sigma}_3$ one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $\beta_2^{-1}(0)$ and $(\beta_1+\beta_2)^{-1}(\pi)$, respectively. Then we have

$$\begin{aligned}X_{\mathbf{x}}^{\widetilde{\sigma}_1}&=-2(2\cot 2x_1+(q-2)\cot x_1)(e_1+e_2),\\ X_{\mathbf{x}}^{\widetilde{\sigma}_2}&=-2(q\cot x_1+\cot 2x_1)e_1,\\ X_{\mathbf{x}}^{\widetilde{\sigma}_3}&=-2(2\cot 2x_1+(q-2)\cot x_1)(e_1-e_2).\end{aligned}$$

Hence the orbit $S(U(2)\times U(q))(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x}\in\widetilde{\sigma}_1$) is minimal if and only if $2\cot 2x_1+(q-2)\cot x_1=0$ ($x_2=x_1$) holds, that is, $(x_1,x_2)=(\arctan\sqrt{q-1},\arctan\sqrt{q-1})$. Hence the orbit $S(U(2)\times U(q))(\mathrm{Exp}(\arctan\sqrt{q-1}(e_1+e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_1$. Also, the orbit $S(U(2)\times U(q))(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x}\in\widetilde{\sigma}_2$) is minimal if and only if $q\cot x_1+\cot 2x_1=0$ ($x_2=0$) holds, that is, $(x_1,x_2)=(\arctan\sqrt{2q+1},0)$. Hence the orbit $S(U(2)\times U(q))(\mathrm{Exp}(\arctan\sqrt{2q+1}e_1))$ is the only minimal singular orbit through $\widetilde{\sigma}_2$. Also, the orbit $S(U(2)\times U(q))(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x}\in\widetilde{\sigma}_3$) is minimal if and only if $2\cot 2x_1+(q-2)\cot x_1=0$ ($x_2=\pi-x_1$) holds, that is, $(x_1,x_2)=(\arctan\sqrt{q-1},\pi-\arctan\sqrt{q-1})$. Hence the orbit $S(U(2)\times U(q))(\mathrm{Exp}(\arctan\sqrt{q-1}e_1+(\pi-\arctan\sqrt{q-1})e_2))$ is the only minimal singular orbit through $\widetilde{\sigma}_3$.



Fig. 7.

Easily we can show $\operatorname{div} X > 0$ (on $\widetilde{C}$). Therefore, since X has the only zero point and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_i\,(i=1,2,3)$ by the proof of Theorem A, all the integral curves of X through points other than its zero point converge to points of $\partial\widetilde{C}$ in finite time. Also, we can show $\operatorname{div} X^{\widetilde{\sigma}_i} > 0$ on $\widetilde{\sigma}_i$ $(i=1,2,3)$. Hence all integral curves of $X^{\widetilde{\sigma}_i}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_i$ in finite time. Therefore we obtain the following fact.

Proposition 4.3. *The mean curvature flow having any non-minimal principal orbit of the $S(U(2)\times U(q))$-action (on $SU(2+q)/S(U(2)\times U(q))$) as initial data converges to some singular orbit in finite time. Also, the mean curvature flow having any non-minimal singular orbit through $\exp^\perp(\widetilde{\sigma}_i)$ of the $S(U(2)\times U(q))$-action as initial data converges to one of two singular orbits through $\exp^\perp(\partial\widetilde{\sigma}_i)$ in finite time.*

Example 4. We consider the isotropy action $S(U(p)\times U(p)) \curvearrowright SU(2p)/S(U(p)\times U(p))$. Then, since $\triangle'$ is the root system of $SU(2p)/S(U(p)\times U(p))$, it is of (C_p)-type and $\mathfrak{b}, \triangle'_+, \Pi$ and δ are described as

$$\begin{aligned}
&\mathfrak{b}=\operatorname{Span}\{e_1,\cdots,e_p\},\\
&\triangle'_+=\{\beta_i-\beta_j\,|\,1\le i<j\le p\}\cup\{\beta_i+\beta_j\,|\,1\le i<j\le p\}\\
&\qquad\cup\{2\beta_i\,|\,1\le i\le p\},\\
&\Pi=\{\beta_i-\beta_{i+1}\,|\,1\le i\le p-1\}\cup\{2\beta_p\},\ \ \delta=2\beta_1,
\end{aligned}\tag{4.19}$$

where $(e_1,\cdots,e_p)$ is the orthonormal base of $\mathfrak{b}$ and $(\beta_1,\cdots,\beta_p)$ is the dual base of

$(e_1, \cdots, e_p)$. For simplicity, we set $\beta_{ij} := \beta_i - \beta_j$ $(1 \leq i < j \leq p)$, $I := \{1, \cdots, p\}$ and $\widehat{I} := \{(i,j) \in I^2 \,|\, 1 \leq i < j \leq p\}$. It is shown that β_{ij}'s $((i,j) \in \hat{I})$ and $\beta_i + \beta_j$'s $((i,j) \in \hat{I})$ are of multiplicity 2 and $2\beta_i$'s $(i \in I)$ are of multiplicity 1. Since we consider the isotropy action, we have $\triangle'^V_+ = \triangle'_+$ and $\triangle'^H_+ = \emptyset$, that is,

$$\begin{aligned}
\widetilde{C} &= \{\mathbf{x} \in \mathfrak{b} \,|\, 0 < \beta_{ij}(\mathbf{x}) < \pi \ ((i,j) \in \widehat{I}),\ 0 < (\beta_i + \beta_j)(\mathbf{x}) < \pi \ ((i,j) \in \widehat{I}), \\
&\qquad\qquad\qquad\qquad\qquad\qquad 0 < 2\beta_i(\mathbf{x}) < \pi \ (i \in I)\} \\
&= \{\mathbf{x} \in \mathfrak{b} \,|\, \beta_{i,i+1}(\mathbf{x}) > 0 \, (1 \leq i \leq p-1),\ 2\beta_p(\mathbf{x}) > 0,\ 2\beta_1(\mathbf{x}) < \pi\}.
\end{aligned}$$

From (4.5), we can describe X explicitly as

$$\begin{aligned}
X_{\mathbf{x}} &= -\sum_{(i,j)\in\widehat{I}} 2\cot(x_i - x_j)(e_i - e_j) - \sum_{(i,j)\in\widehat{I}} 2\cot(x_i + x_j)(e_i + e_j) \\
&\quad - \sum_{i\in I} \cot 2x_i \cdot 2e_i \\
&= -2\sum_{i\in I}\left(\sum_{j\in I\setminus\{i\}} (\cot(x_i - x_j) + \cot(x_i + x_j)) + \cot 2x_i\right) e_i.
\end{aligned}$$

According to this relation, a principal orbit $S(U(p) \times U(p))(\operatorname{Exp}\mathbf{x})$ $(\mathbf{x} \in \widetilde{C})$ is minimal if and only if the following relations hold:

$$\sum_{j\in I\setminus\{i\}} (\cot(x_i - x_j) + \cot(x_i + x_j)) + \cot 2x_i = 0 \quad (i \in I). \tag{4.20}$$

In the sequel, we consider the case of $p = 2$. Then (4.20) is as follows:

$$\left\{\begin{array}{l} \cot(x_1 - x_2) + \cot(x_1 + x_2) + \cot 2x_1 = 0 \\ \cot(x_2 - x_1) + \cot(x_1 + x_2) + \cot 2x_2 = 0. \end{array}\right. \tag{4.21}$$

Under $0 < x_1 - x_2 < \pi$, $0 < x_1 + x_2 < \pi$, $0 < 2x_1 < \pi$ and $0 < 2x_2 < \pi$, the equation (4.21) has the only solution, which we denote by (α_1, α_2). Therefore the orbit $S(U(2) \times U(2))(\operatorname{Exp}(\alpha_1 e_1 + \alpha_2 e_2))$ is the only minimal principal orbit of the $S(U(2) \times U(2))$-action. Denote by $\widetilde{\sigma}_1$, $\widetilde{\sigma}_2$ and $\widetilde{\sigma}_3$ the one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $(2\beta_2)^{-1}(0)$ and $(2\beta_1)^{-1}(\pi)$. Then we have

$$\begin{aligned}
X_{\mathbf{x}}^{\widetilde{\sigma}_1} &= -4\cot 2x_1(e_1 + e_2), \\
X_{\mathbf{x}}^{\widetilde{\sigma}_2} &= -2(2\cot x_1 + \cot 2x_1)e_1, \\
X_{\mathbf{x}}^{\widetilde{\sigma}_3} &= -2\cot 2x_2 e_2.
\end{aligned}$$

Hence the orbit $S(U(2) \times U(2))(\operatorname{Exp}\mathbf{x})$ $(\mathbf{x} \in \widetilde{\sigma}_1)$ is minimal if and only if $\cot 2x_1 = 0$ $(x_2 = x_1)$ holds, that is, $(x_1, x_2) = (\frac{\pi}{4}, \frac{\pi}{4})$. Therefore the orbit $S(U(2) \times U(2))$

$(\mathrm{Exp}(\frac{\pi}{4}(e_1+e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_1$. Also, the orbit $S(U(2)\times U(2))(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x}\in\widetilde{\sigma}_2)$ is minimal if and only if $2\cot x_1+\cot 2x_1=0$ $(x_2=0)$ holds, that is, $(x_1,x_2)=(\arctan\sqrt{5},0)$. Therefore the orbit $S(U(2)\times U(2))(\mathrm{Exp}(\arctan\sqrt{5}e_1))$ is the only minimal singular orbit through $\widetilde{\sigma}_2$. Also, the orbit $S(U(2)\times U(2))(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x}\in\widetilde{\sigma}_3)$ is minimal if and only if $\cot 2x_2=0$ $(x_1=\frac{\pi}{2})$ holds, that is, $(x_1,x_2)=(\frac{\pi}{2},\frac{\pi}{4})$. Therefore the orbit $S(U(2)\times U(2))(\mathrm{Exp}(\frac{\pi}{4}(2e_1+e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_3$.



Fig. 8.

Easily we can show $\mathrm{div}\,X>0$ (on $\widetilde{C}$). Therefore, since X has the only zero point and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_i\,(i=1,2,3)$ by the proof of Theorem A, all the integral curves of X through points other than its zero point converge to points of $\partial\widetilde{C}$ in finite time. Also, we can show $\mathrm{div}\,X^{\widetilde{\sigma}_i}>0$ on $\widetilde{\sigma}_i$ $(i=1,2,3)$. Hence all integral curves of $X^{\widetilde{\sigma}_i}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_i$ in finite time. Therefore we obtain the following fact.

Proposition 4.4. *The mean curvature flow having any non-minimal principal orbit of the* $S(U(2)\times U(2))$*-action (on* $SU(4)/S(U(2)\times U(2))$*) as initial data converges to some singular orbit in finite time. Also, the mean curvature flow having any non-minimal singular orbit through* $\exp^\perp(\widetilde{\sigma}_i)$ *of the* $S(U(2)\times U(2))$*-action as initial data converges to one of two singular orbits through* $\exp^\perp(\partial\widetilde{\sigma}_i)$ *in finite time.*

Example 5. We consider the Hermann action $SO(p+q) \curvearrowright SU(p+q)/S(U(p)\times U(q))$ $(p<q)$. Since $L/H\cap K$ is then equal to $SO(p+q)/SO(p)\times SO(q)$, the cohomogeneity of this action is equal to p, that is, it is the rank of $SU(p+q)/S(U(p)\times U(q))$. Hence a maximal abelian subspace $\mathfrak{b}$ of $\mathfrak{p}\cap\mathfrak{q}$ is also a maximal abelian subspace of $\mathfrak{p}$ and hence $\triangle'$ is the root system of $SU(p+q)/S(U(p)\times U(q))$. Hence $\triangle'$ is of (B_p)-type and the quantities $\mathfrak{b}$, $\triangle'_+$, Π and δ are as in (4.16). Let β_i, β_{ij}, I and $\widehat{I}$ be as in Example 3. We have $\dim\mathfrak{p}_{\beta_{ij}}=\dim\mathfrak{p}_{\beta_i+\beta_j}=2\,((i,j)\in\hat{I})$, $\dim\mathfrak{p}_{\beta_i}=2(q-p)\,(i\in I)$ and $\dim\mathfrak{p}_{2\beta_i}=1\,(i\in I)$. Also, since ${\triangle'}^V$ is the root system of $SO(p+q)/SO(p)\times SO(q)$, it is of (B_p)-type. The Satake diagram of $SO(p+q)/SO(p)\times SO(q)$ is as follows:



$(p+q\,:\,\text{odd})$

$(p+q\,:\,\text{even},\;q-p\geq 4)$

$(p+q\,:\,\text{even},\;q-p=2).$

Hence we have ${\triangle'}^V_+=\triangle'_+\setminus\{2\beta_i\,|\,i\in I\}$, $\dim(\mathfrak{p}_{\beta_{ij}}\cap\mathfrak{q})=\dim(\mathfrak{p}_{\beta_i+\beta_j}\cap\mathfrak{q})=1$ $((i,j)\in\hat{I})$ and $\dim(\mathfrak{p}_{\beta_i}\cap\mathfrak{q})=q-p$ $(i\in I)$. Furthermore we have ${\triangle'}^H_+=\triangle'_+$, $\dim(\mathfrak{p}_{\beta_{ij}}\cap\mathfrak{h})=\dim(\mathfrak{p}_{\beta_i+\beta_j}\cap\mathfrak{h})=1$ $((i,j)\in\hat{I})$, $\dim(\mathfrak{p}_{\beta_i}\cap\mathfrak{h})=q-p$ $(i\in I)$ and $\dim(\mathfrak{p}_{2\beta_i}\cap\mathfrak{h})=1$ $(i\in I)$. Therefore we have

$$\widetilde{C}=\{\mathbf{x}\in\mathfrak{b}\,|\,\beta_{i,i+1}(\mathbf{x})>0\ (i=1,\cdots,p-1),\ \beta_p(\mathbf{x})>0,\ (\beta_1+\beta_2)(\mathbf{x})<\frac{\pi}{2}\}.$$

From (4.5), we can describe X explicitly as

$$\begin{aligned}X_{\mathbf{x}}=&\sum_{i\in I}2\tan 2x_i\cdot e_i-\sum_{(i,j)\in\widehat{I}}2\cot 2(x_i-x_j)(e_i-e_j)\\&-\sum_{i\in I}2(q-p)\cot 2x_i\cdot e_i-\sum_{(i,j)\in\widehat{I}}2\cot 2(x_i+x_j)(e_i+e_j)\\=&\,2\sum_{i\in I}\left(\tan 2x_i-(q-p)\cot 2x_i-\sum_{i\in I\setminus\{i\}}(\cot 2(x_i-x_j)+\cot 2(x_i+x_j))\right)e_i.\end{aligned}$$

According to this relation, a principal orbit $SO(p+q)(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x}\in\widetilde{C})$ is minimal if and only if the following relations hold:

$$\tan 2x_i-(q-p)\cot 2x_i-\sum_{j\in I\setminus\{i\}}(\cot 2(x_i-x_j)+\cot 2(x_i+x_j))=0\quad(i\in I).\tag{4.22}$$

In the sequel, we consider the case of $p=2$. Then (4.22) is as follows:

$$\left\{\begin{array}{l}\tan 2x_1-(q-2)\cot 2x_1-\cot 2(x_1-x_2)-\cot 2(x_1+x_2)=0\\\tan 2x_2-(q-2)\cot 2x_2-\cot 2(x_2-x_1)-\cot 2(x_1+x_2)=0.\end{array}\right.\tag{4.23}$$

Under $0 < x_1 - x_2 < \frac{\pi}{2}$, $0 < x_1 + x_2 < \frac{\pi}{2}$, $0 < x_i < \frac{\pi}{2}\,(i = 1, 2)$ and $-\frac{\pi}{2} < 2x_i < \frac{\pi}{2}\,(i = 1, 2)$, the equation (4.23) has the only solution, which we denote by (α_1, α_2). Here we note that $\lim\limits_{q\to\infty}(\alpha_1, \alpha_2) = (\frac{\pi}{4}, \frac{\pi}{4})$. Denote by $\widetilde{\sigma}_1$, $\widetilde{\sigma}_2$ and $\widetilde{\sigma}_3$ the one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $\beta_2^{-1}(0)$ and $(\beta_1 + \beta_2)^{-1}(\frac{\pi}{2})$, respectively. Then we have

$$\begin{aligned}
X_{\mathbf{x}}^{\widetilde{\sigma}_1} &= 2(\tan 2x_1 - (q-2)\cot 2x_1 - \cot 4x_1)(e_1 + e_2),\\
X_{\mathbf{x}}^{\widetilde{\sigma}_2} &= 2(\tan 2x_1 - q\cot 2x_1)e_1,\\
X_{\mathbf{x}}^{\widetilde{\sigma}_3} &= 2(\tan 2x_1 - (q-2)\cot 2x_1 - \cot 4x_1)(e_1 - e_2).
\end{aligned}$$

Hence the orbit $SO(q+2)(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x} \in \widetilde{\sigma}_1)$ is minimal if and only if $\tan 2x_1 - (q-2)\cot 2x_1 - \cot 4x_1 = 0$ $(x_2 = x_1)$ holds, that is, $(x_1, x_2) = (\frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1}, \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1})$. Hence the orbit $SO(q+2)(\mathrm{Exp}(\frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1}(e_1+e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_1$. Also, the orbit $SO(q+2)(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x} \in \widetilde{\sigma}_2)$ is minimal if and only if $\tan 2x_1 - q\cot 2x_1 = 0$ $(x_2 = 0)$ holds, that is, $(x_1, x_2) = (\frac{1}{2}\arctan\sqrt{q}, 0)$. Hence the orbit $SO(q+2)(\mathrm{Exp}(\frac{1}{2}\arctan\sqrt{q}e_1))$ is the only minimal singular orbit through $\widetilde{\sigma}_2$. Also, the orbit $SO(q+2)(\mathrm{Exp}\,\mathbf{x})$ $(\mathbf{x} \in \widetilde{\sigma}_3)$ is minimal if and only if $\tan 2x_1 - (q-2)\cot 2x_1 - \cot 4x_1 = 0$ $(x_2 = \frac{\pi}{2} - x_1)$ holds, that is, $(x_1, x_2) = (\frac{\pi}{2} - \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1}, \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1})$. Hence the orbit $SO(q+2)(\mathrm{Exp}((\frac{\pi}{2} - \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1})e_1 + \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1e_2})$ is the only minimal singular orbit through $\widetilde{\sigma}_3$.


$\beta_1^{-1}(\frac{\pi}{2})$
$\beta_{12}^{-1}(\frac{\pi}{2})$
$(\beta_1 + \beta_2)^{-1}(\frac{\pi}{2})$
$\alpha_1 e_1 + \alpha_2 e_2$
$\frac{\pi}{4}e_1$
$\frac{1}{2}\arctan\sqrt{q}e_1$
$(\frac{\pi}{2} - \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1})e_1 + \frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1e_2}$
$\beta_2^{-1}(0)$
$(q \to \infty)$
$(q \to \infty)$
$\beta_1^{-1}(0)$
$(q \to \infty)$
$\mathbf{0}$
$(q \to \infty)$
$\beta_{12}^{-1}(0)$
$\frac{1}{2}\arctan\sqrt{\frac{2}{3}q - 1}(e_1 + e_2)$
$\beta_2^{-1}(\frac{\pi}{4})$

Fig. 9.

Easily we can show $\operatorname{div} X > 0$ (on $\widetilde{C}$). Therefore, since X has the only zero point and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_i$ $(i = 1, 2, 3)$ by the proof of Theorem A, all the integral curves of X through points other than its zero point converge to points of $\partial\widetilde{C}$ in finite time. Also, we can show $\operatorname{div} X^{\widetilde{\sigma}_i} > 0$ on $\widetilde{\sigma}_i$ $(i = 1, 2, 3)$. Hence all integral curves of $X^{\widetilde{\sigma}_i}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_i$ in finite time. Therefore we obtain the following fact.

Proposition 4.5. *The mean curvature flow having any non-minimal principal orbit of the $SO(q+2)$-action (on $SU(q+2)/S(U(2)\times U(q))$) as initial data converges to some singular orbit in finite time, where $q > 2$. Also, the mean curvature flow having any non-minimal singular orbit through $\exp^{\perp}(\widetilde{\sigma}_i)$ of the $SO(q+2)$-action as initial data converges to one of two singular orbits through $\exp^{\perp}(\partial\widetilde{\sigma}_i)$ in finite time.*

Example 6. We consider the Hermann action $SO(2p) \curvearrowright SU(2p)/S(U(p)\times U(p))$. Since $L/H\cap K$ is then equal to $SO(2p)/SO(p)\times SO(p)$, the cohomogeneity of this action is equal to $p\,(=\operatorname{rank}(SU(2p)/S(U(p)\times U(p))))$. Hence a maximal abelian subspace $\mathfrak{b}$ of $\mathfrak{p}\cap\mathfrak{q}$ is also a maximal abelian subspace of $\mathfrak{p}$ and hence $\triangle'$ is the root system of $SU(2p)/S(U(p)\times U(p))$. Hence it is of (C_p)-type and the quantities $\mathfrak{b}$, $\triangle'_+$, Π and δ are as in (4.19). Let β_i, β_{ij}, I and $\widehat{I}$ be as in Example 4. We have $\dim\mathfrak{p}_{\beta_{ij}} = \dim\mathfrak{p}_{\beta_i+\beta_j} = 2$ $((i,j)\in\hat{I})$ and $\dim\mathfrak{p}_{2\beta_i} = 1$ $(i\in I)$. Also, ${\triangle'}^V$ is the root system of $SO(2p)/SO(p)\times SO(p)$. Hence it is of (D_p)-type and we have ${\triangle'}^V_+ = \{\beta_{ij}\,|\,(i,j)\in\hat{I}\}\cup\{\beta_i+\beta_j\,|\,(i,j)\in\hat{I}\}$ and $\dim(\mathfrak{p}_{\beta_{ij}}\cap\mathfrak{q}) = \dim(\mathfrak{p}_{\beta_i+\beta_j}\cap\mathfrak{q}) = 1$ $((i,j)\in\hat{I})$. Hence we have ${\triangle'}^H_+ = \triangle'_+$ and each root of ${\triangle'}^H_+$ has multiplicity 1. Therefore we have

$$\widetilde{C} = \{\mathbf{x}\in\mathfrak{b}\,|\,\beta_{i,i+1}(\mathbf{x}) > 0\ (i = 1,\cdots,p-1),\ (\beta_{p-1}+\beta_p)(\mathbf{x}) > 0,\ 2\beta_1(\mathbf{x}) < \frac{\pi}{2}\}$$

and

$$\begin{aligned} X_{\mathbf{x}} &= -\sum_{(i,j)\in\widehat{I}} 2\left(\cot 2(x_i-x_j)(e_i-e_j) + \cot 2(x_i+x_j)(e_i+e_j)\right) \\ &\quad + \sum_{i\in I}\tan 2x_i\cdot 2e_i \\ &= 2\sum_{i\in I}\left(-\sum_{j\in I\setminus\{i\}}(\cot 2(x_i-x_j)+\cot 2(x_i+x_j)) + \tan 2x_i\right)e_i. \end{aligned}$$

According to this relation, a principal orbit $SO(2p)(\operatorname{Exp}\mathbf{x})$ $(\mathbf{x}\in\widetilde{C})$ is minimal if and only if the following relation holds:

$$\sum_{j\in I\setminus\{i\}}(\cot 2(x_i-x_j)+\cot 2(x_i+x_j)) - \tan 2x_i = 0 \quad (i\in I). \tag{4.24}$$

In the sequel, we consider the case of $p = 2$. Then (4.24) is as follows:

$$\left\{\begin{array}{l}\cot 2(x_1 - x_2) + \cot 2(x_1 + x_2) - \tan 2x_1 = 0\\ \cot 2(x_2 - x_1) + \cot 2(x_1 + x_2) - \tan 2x_2 = 0.\end{array}\right. \tag{4.25}$$

Under $x_1 - x_2 > 0$, $x_1 + x_2 > 0$ and $x_1 < \frac{\pi}{4}$, the equation (4.25) has the only solution $(x_1, x_2) = (\frac{1}{2}\arctan\sqrt{2}, 0)$. Therefore the orbit $SO(4)\mathrm{Exp}(\frac{1}{2}\arctan\sqrt{2}e_1)$ is the only minmal principal orbit of the $SO(4)$-action. Denote by $\widetilde{\sigma}_1$, $\widetilde{\sigma}_2$ and $\widetilde{\sigma}_3$ the one dimensional stratums of $\partial\widetilde{C}$ which are contained in $\beta_{12}^{-1}(0)$, $(\beta_1+\beta_2)^{-1}(0)$ and $(2\beta_1)^{-1}(\frac{\pi}{2})$, respectively. Then we have

$$\begin{aligned}
X_{\mathbf{x}}^{\widetilde{\sigma}_1} &= -2(\cot 2(x_1 + x_2) - \tan 2x_1)e_1 - 2(\cot 2(x_1 + x_2) - \tan 2x_2)e_2\\
X_{\mathbf{x}}^{\widetilde{\sigma}_2} &= -2(\cot 2(x_1 - x_2) - \tan 2x_1)e_1 - 2(\cot 2(x_2 - x_1) - \tan 2x_2)e_2\\
X_{\mathbf{x}}^{\widetilde{\sigma}_3} &= -2(\cot 2(x_1 - x_2) + \cot 2(x_1 + x_2))e_1\\
&\qquad -2(\cot 2(x_2 - x_1) + \cot 2(x_1 + x_2) - \tan 2x_2)e_2.
\end{aligned}$$

Hence the orbit $SO(4)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{\sigma}_1$) is minimal if and only if $\cot 4x_1 = \tan 2x_1$ ($x_2 = x_1$) holds, that is, $(x_1, x_2) = (\frac{\pi}{12}, \frac{\pi}{12})$. Therefore the orbit $SO(4)(\mathrm{Exp}(\frac{\pi}{12}(e_1 + e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_1$. Also, the orbit $SO(4)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{\sigma}_2$) is minimal if and only if $\cot 4x_1 = \tan 2x_1$ ($x_2 = -x_1$) holds, that is, $(x_1, x_2) = (\frac{\pi}{12}, -\frac{\pi}{12})$. Therefore the orbit $SO(4)(\mathrm{Exp}(\frac{\pi}{12}(e_1 - e_2)))$ is the only minimal singular orbit through $\widetilde{\sigma}_2$. Also, the orbit $SO(4)(\mathrm{Exp}\,\mathbf{x})$ ($\mathbf{x} \in \widetilde{\sigma}_3$) is minimal if and only if $\cot(\frac{\pi}{2} - 2x_2) + \cot(\frac{\pi}{2} + 2x_2) = 0$ and $-\cot(\frac{\pi}{2} - 2x_2) + \cot(\frac{\pi}{2} + 2x_2) - \tan 2x_2 = 0$ ($x_1 = \frac{\pi}{4}$) hold, that is, $(x_1, x_2) = (\frac{\pi}{4}, 0)$. Therefore the orbit $SO(4)(\mathrm{Exp}(\frac{\pi}{4}e_1))$ is the only minimal singular orbit through $\widetilde{\sigma}_3$.


$(\beta_1+\beta_2)^{-1}(\frac{\pi}{2})$
$\beta_{12}^{-1}(0)$
$\frac{\pi}{4}e_1$
$\frac{1}{2}\arctan\sqrt{2}e_1$
$(2\beta_1)^{-1}(\frac{\pi}{2})$
$\frac{\pi}{12}(e_1+e_2)$
$\beta_{12}^{-1}(\frac{\pi}{2})$
$\mathbf{0}$
$(\beta_1+\beta_2)^{-1}(0)$
$\frac{\pi}{12}(e_1-e_2)$

Fig. 10.

Easily we can show $\operatorname{div} X > 0$ (on $\widetilde{C}$). Therefore, since X has the only zero point and X is as in Fig.1 over a collar neighborhood of $\widetilde{\sigma}_i\,(i = 1, 2, 3)$ by the proof of Theorem A, all the integral curves of X through points other than its zero point converge to points of $\partial\widetilde{C}$ in finite time. Also, we can show $\operatorname{div} X^{\widetilde{\sigma}_i} > 0$ on $\widetilde{\sigma}_i$ $(i = 1, 2, 3)$. Hence all integral curves of $X^{\widetilde{\sigma}_i}$ through points other than its zero point converge to points of $\partial\widetilde{\sigma}_i$ in finite time. Therefore we obtain the following fact.

Proposition 4.6. *The mean curvature flow having any non-minimal principal orbit of the $SO(4)$-action (on $SU(4)/S(U(2) \times U(2))$) as initial data converges to some singular orbit in finite time. Also, the mean curvature flow having any non-minimal singular orbit through $\exp^{\perp}(\widetilde{\sigma}_i)$ of the $SO(4)$-action as initial data converges to one of two singular orbits through $\exp^{\perp}(\partial\widetilde{\sigma}_i)$ in finite time.*

References

[BV] J. Berndt and L. Vanhecke, Curvature-adapted submanifolds, Nihonkai Math. J. **3** (1992) 177-185.

[BCO] J. Berndt, S. Console and C. Olmos, Submanifolds and holonomy, Research Notes in Mathematics 434, CHAPMAN & HALL/CRC Press, Boca Raton, London, New York Washington, 2003.

[Ch] U. Christ, Homogeneity of equifocal submanifolds, J. Differential Geometry **62** (2002) 1-15.

[Co] L. Conlon, Remarks on commuting involutions, Proc. Amer. Math. Soc. **22** (1969) 255-257.

[GT] O. Goertsches and G. Thorbergsson, On the Geometry of the orbits of Hermann actions, Geom. Dedicata **129** (2007) 101-118.

[H] S. Helgason, Differential geometry, Lie groups and symmetric spaces, Academic Press, New York, 1978.

[HLO] E, Heintze, X. Liu and C. Olmos, Isoparametric submanifolds and a Chevalley-type restriction theorem, Integrable systems, geometry, and topology, 151-190, AMS/IP Stud. Adv. Math. 36, Amer. Math. Soc., Providence, RI, 2006.

[HPTT] E. Heintze, R.S. Palais, C.L. Terng and G. Thorbergsson, Hyperpolar actions on symmetric spaces, Geometry, topology and physics for Raoul Bott (ed. S. T. Yau), Conf. Proc. Lecture Notes Geom. Topology **4**, Internat. Press, Cambridge, MA, 1995 pp214-245.

[Koi1] N. Koike, On proper Fredholm submanifolds in a Hilbert space arising from submanifolds in a symmetric space, Japan. J. Math. **28** (2002) 61-80.

[Koi2] N. Koike, Actions of Hermann type and proper complex equifocal submanifolds, Osaka J. Math. **42** (2005) 599-611.

[Kol] A. Kollross, A Classification of hyperpolar and cohomogeneity one actions, Trans.

Amer. Math. Soc. **354** (2001) 571-612.
[LT] X. Liu and C. L. Terng, The mean curvature flow for isoparametric submanifolds, Duke Math. J. **147** (2009) 157-179.
[P] R.S. Palais, Morse theory on Hilbert manifolds, Topology **2** (1963) 299-340.
[PT] R.S. Palais and C.L. Terng, Critical point theory and submanifold geometry, Lecture Notes in Math. **1353**, Springer, Berlin, 1988.
[T1] C.L. Terng, Isoparametric submanifolds and their Coxeter groups, J. Differential Geometry **21** (1985) 79-107.
[T2] C.L. Terng, Proper Fredholm submanifolds of Hilbert space, J. Differential Geometry **29** (1989) 9-47.
[TT] C.L. Terng and G. Thorbergsson, Submanifold geometry in symmetric spaces, J. Differential Geometry **42** (1995) 665-718.
[Z] X. P. Zhu, Lectures on mean curvature flows, Studies in Advanced Math., AMS/IP, 2002.

Department of Mathematics, Faculty of Science
Tokyo University of Science, 26 Wakamiya-cho
Shinjuku-ku, Tokyo 162-8601 Japan
(koike@ma.kagu.tus.ac.jp)